
1 - Murray River & Floodplain –Merbein to SA Border

Regional Catchment Strategy Implementation Plan



Mallee Regional Catchment Strategy 2013-19

This plan is a living document, which may be updated as required in light of new information or changing conditions. Whenever this document is revised, it will be uploaded to the Mallee CMA Website, replacing the old version.

To allow for ongoing review and renewal processes, some sections of the document may not be entirely complete at the time of publishing.

Publications produced by the Mallee Catchment Management Authority may be of assistance to you but the Mallee Catchment Management Authority and its employees do not guarantee that the publication is without flaw of any kind or is wholly appropriate for your particular purpose and therefore disclaims all liability for any error, loss or other consequence which may arise from you relying on any information in any Mallee Catchment Management Authority publication.

Version Control						
Version Number	Prepared by	Reviewed by	Date			
1 (Presented on internal blog)	Sean Dwyer	Jo Latta	06/11/2013			
2	Amy Leamon	Jo Latta	17/03/2017			

[©] Mallee Catchment Management Authority 2017

Contents

Section 1 - Overview	5
Catchment Asset Significance	6
Catchment Asset Value	6
Threatening Processes	7
Management Plans	8
Management Actions	8
Priorities	8
Regional Delivery Partners	11
Community NRM Groups	11
Section 2 – Regional Assets in this Catchment Asset	14
Rivers	15
Wetlands	16
Threatened Species and Communities	16
Terrestrial Habitat	23
Soils	26
Agricultural Land	27
Groundwater	28
Cultural Heritage	28
Community Capacity	29
Section 3 – Threatening Processes in this Catchment Asset	32
Land & water salinisation	33
Invasive Plants	35
Invasive Animals	37
Altered Hydrological Regimes	39
Soil Erosion	40
Inappropriate Water Use Practices	42
Recreational Pressures	43
Land Use Change	43
Direct Off-site Interactions	44
Misaligned Community Perceptions	44
Inappropriate Fire Regimes	45
Constrained Regenerative Capacity	46
Section 4 – References & Appendices	48
References	49
The Value of Regional Assets within a Catchment Asset	50

2. Categorising Threatening Processes	50
3. Management Plans Relevant to this Catchment Asset	51
4. Management Action Definitions	54
5. Management Actions from Existing Management Plans	55
6. Prioritising Potential Management Actions	63
7. Management Action Priorities	64

Section 1 - Overview

Catchment Asset Significance

The Merbein to SA Border reach of the Murray River begins at Merbein and extends approximately 232 kilometres downstream to the South Australian Border. The area forms the northern part of the region commonly referred to as the Millewa. It occupies an area of approximately 50,000 hectares that is predominantly given over to native vegetation reserved on public land.

This area is important for its high diversity of both terrestrial and aquatic habitats that provide for significant populations of some rare and threatened species and ecological communities. These include the Regent Parrot (Polytelis anthopeplus monarchoides), Giles Planingale (Planigale gilesi), Winged Peppercress (Lepidium monoplocoides) and the Lowland Riverine Fish Community.

Within this Catchment Asset, there are two nationally recognised Living Murray Icon Sites: the Murray River Channel; and part of the Chowilla Floodplains Living Murray Icon Site (Lindsay and Wallpolla Islands). Lindsay and Wallpolla Islands and Lake Wallawalla are listed in the Australian Directory of Important Wetlands and they are known to play host to migratory bird species identified in international migratory bird agreements between Australia and Japan (JAMBA), China (CAMBA) and South Korea (ROKAMBA).

The area includes parts of Murray Sunset National Park, which is listed in Category II (National Parks) of the International Union for Conservation of Nature (IUCN) List of National Parks and Protected Areas, and the River Murray Reserve.

The area is of geomorphological significance as Lindsay Island and its anabranch, the Lindsay River, form an excellent example of a permanent river course change produced by a major flood event. The Lindsay River is one of the largest anabranches of the Murray in Victoria.

The asset area provides significant ecological linkages to the River Red Gum Forests of NSW, the Chowilla Floodplains of NSW and South Australia, and Murray-Sunset National Park.

The major land uses in this area are rangelands grazing, forestry, conservation and recreation, with extensive areas of public land, such as the Lindsay Island section of the Murray-Sunset National Park, Wallpolla Island State Forest and the Murray River Public Purposes Reserve.

The area is sparsely populated and isolated from major towns. Intensive recreational use at key times of the year brings thousands of visitors to the area. Recreation is an important use of frontages in the Merbein to SA Border region. Lindsay, Mulcra and Wallpolla Islands are favoured by locals and visitors alike for the camping spots, fishing, canoeing, bird watching, and four-wheel driving.

This area's cultural heritage is of national and international importance. Archaeological evidence shows that Aboriginal people inhabited the surrounding floodplains of the Murray River for thousands of years, on a relatively permanent basis. Throughout this entire area the large concentration of burials is very unique to Victoria. There are also a significantly a large amount of meeting places, ceremonial places, scared trees, and artefact sites throughout this area.

The area also represents early adoption of irrigation in Victoria, for example at Yelta and subsequently in later years, the beginning of the Australian Irrigated Colonies.

Catchment Asset Value

This section examines the relative values of the Regional Assets that lie within this Catchment Asset. It provides an overview of the asset analysis in order to inform prioritisation of management actions that are intended to minimise the impact of the threatening processes upon Regional Assets and improve the status or quality of the Regional Assets within the Catchment Asset Area.

The asset value indicates the relative importance of the Regional Asset within the Catchment Asset Area. The valuation is determined from a range of indicators, many specific to each of the Regional Assets that describe the importance of the Asset relative to social, economic and environmental values.

Table 1 below provides a summary of the value of each of the Regional Assets in this Catchment Asset. Greater detail about each of the Regional Assets in this Catchment Asset can be found in Section 2 – Regional Assets in this Catchment Asset

Table 1: Value of Regional Assets in this Catchment Asset

Regional Asset	Value
Rivers	VERY HIGH
Wetlands	VERY HIGH
Threatened Species and Communities	HIGH
Terrestrial Habitat	VERY HIGH
Soils	HIGH
Agricultural Land	HIGH
Groundwater	NONE
Cultural Heritage	HIGH
Community Capacity	HIGH

A detailed explanation of how these values were defined and applied can be found in Appendix 1. The Value of Regional Assets within a Catchment Asset.

Threatening Processes

This section examines the threatening processes that may be impacting upon Regional Assets that lie within this Catchment Asset. It provides an overview of each of the threatening processes in order to inform prioritisation of management actions that are intended to minimise their impact upon Regional Assets and improve the status or quality of the Regional Assets within the Catchment Asset Area.

Table 2 below provides a summary of the threat level posed by each of the threatening processes that are active in this Catchment Asset. Greater detail about each of the threatening processes and their scope, scale and relative impact within this Catchment Asset can be found in Section 3 – Threatening Processes in this Catchment Asset.

Table 2: Threat Priority across this Catchment Asset

Threatening Process	Priority of Action
Land & Water Salinisation	MEDIUM
Invasive Plants	HIGH
Invasive Animals	HIGH
Altered Hydrological Regimes	HIGH
Soil Erosion	HIGH
Inappropriate Water Use Practices	HIGH
Recreational Pressures	HIGH
Land Use Change	HIGH
Direct off-site interactions	LOW

Misaligned community perceptions	MEDIUM
Inappropriate fire regimes	MEDIUM
Constrained regenerative capacity	HIGH

A detailed explanation of how these threat levels were defined and applied can be found in Appendix 2. Categorising Threatening Processes.

Management Plans

This Catchment Asset Area is subject to a diverse range of natural resource management instruments that are intended to protect, preserve and enhance the area's natural resources for the benefit of the Mallee region. These instruments can be either broad in their focus (covering many regional assets), focused on a particular area of interest (such as an individual Regional Asset) or tightly focused on a specific location or species. These management plans are prepared and owned by range of entities such as local government, statutory authorities, community groups and government departments at both State and Federal levels. Responsibility for their implementation can rest with a single entity or it may be divested across a broad range of organisations and groups. Some management plans are prepared to satisfy a legislative requirement while others are prepared to provide direction towards an organisational goal. Regardless of the reason and purpose of the management plan, they all have a common feature – they typically contain specific management direction or actions that are to be delivered through the implementation of the plan.

The identified existing management plans have been listed in Appendix 3. *Management Plans Relevant to this Catchment Asset*

Management Actions

Actions within the management plans above are typically diverse in their wording, structure and complexity. However, they are broadly similar in terms of their intent. As a result, they can usually be easily classified into broad categories. Within each of these categories are a range of common management actions. Each of these common management actions can also be said to target specific threatening processes. These management categories, the management actions within each of them, their definitions and the targeted threatening processes are shown in Appendix 4. *Management Action Definitions*

The result of the classification process of the actions within Appendix 3. Management Plans Relevant to this Catchment Asset is contained in Appendix 5. Management Actions from Existing Management Plans. The classification table includes some detail about each action to support the classification. It is advised that if more information about a specific management action within a management plan is required then the original management plan should be consulted rather than relying specifically on the data in that classification table.

Priorities

This section examines the potential management action groups from the management plans relevant to this Catchment Asset Area that are listed in the Management Plans section above and summarised Appendix 5. Management Actions from Existing Management Plans. The intent is to prioritise these potential management actions in terms of:

- the priority of the regional asset within the catchment asset area (outlined in the Catchment Asset Value section above);
- the priority to address a threatening process (outlined in the Threatening Processes section above);

- the capacity of the management action to address the threatening process; and
- the level of investment (in both of time and money) required to exercise that capacity.

The individual management actions from the plans listed in the Management Plans section were grouped and scored by consensus against agreed criteria and then each potential management action group within the Catchment Asset was assigned to a category that defines the prospective priority to the implementation. Further information regarding the definition and application of each of these categories can be found in Appendix 6. *Prioritising Potential Management Actions*.

The findings of the analysis are summarised in Appendix 7. Management Action Priorities.

Table 3 below. The detailed priority matrices from which the summary table is taken can be found in Appendix 7. *Management Action Priorities*.

Table 3: Summary of the Priority of Potential Management Actions

Management Action	Priority	Threatening Processes Addressed			
Pest Plant Control	High	Invasive Plants; Constrained Regenerative Capacity			
Pest Animal Control	High	Invasive Animals; Soil Erosion; Constrained Regenerative Capacity			
Habitat Protection	High	Invasive Animals; Soil Erosion; Recreational Pressure; Land Use Change; Direct Off-Site Interactions; Constrained Regenerative Capacity			
Habitat Restoration	Medium	Land & Water Salinisation; Soil Erosion; Recreational Pressure; Direct Off-Site Interactions; Constrained Regenerative Capacity			
Revegetation	Low	Land & Water Salinisation; Soil Erosion; Recreational Pressure; Direct Off-Site Interactions; Constrained Regenerative Capacity			
Environmental Watering	High	Altered Hydrological Regimes; Constrained Regenerative Capacity			
Soil Erosion Control	Medium	Soil Erosion			
Threatened Species Interventions	High	Constrained Regenerative Capacity			
Enhancing Land Management Regimes	High	Land & Water Salinisation; Soil Erosion; Inappropriate Water Use Practises; Land Use Change; Direct Off-Site Interactions			
Supporting Human Capacity for NRM	Medium	Land & Water Salinisation; Invasive Plants; Invasive Animals; Altered Hydrological Regimes; Soil Erosion; Inappropriate Water Use Practices; Recreational Pressures; Land Use Change; Direct Offsite Interactions; Misaligned Community Perceptions; Inappropriate Fire Regimes; Constrained Regenerative Capacity			
Supporting Institutional Capacity for NRM	Medium	Land & Water Salinisation; Invasive Plants; Invasive Animals; Altered Hydrological Regimes; Soil Erosion; Inappropriate Water Use Practices; Recreational Pressures; Land Use Change; Direct Offsite Interactions; Misaligned Community Perceptions; Inappropriate Fire			

		Regimes; Constrained Regenerative Capacity
Supporting Social Capacity for NRM	Medium	Land & Water Salinisation; Invasive Plants; Invasive Animals; Altered Hydrological Regimes; Soil Erosion; Inappropriate Water Use Practices; Recreational Pressures; Land Use Change; Direct Offsite Interactions; Misaligned Community Perceptions; Inappropriate Fire Regimes; Constrained Regenerative Capacity
Institutional Planning for NRM	Medium	Land & Water Salinisation; Invasive Plants; Invasive Animals; Altered Hydrological Regimes; Soil Erosion; Inappropriate Water Use Practices; Recreational Pressures; Land Use Change; Direct Offsite Interactions; Misaligned Community Perceptions; Inappropriate Fire Regimes; Constrained Regenerative Capacity
Community Driven Planning for NRM	Medium	Land & Water Salinisation; Invasive Plants; Invasive Animals; Altered Hydrological Regimes; Soil Erosion; Inappropriate Water Use Practices; Recreational Pressures; Land Use Change; Direct Offsite Interactions; Misaligned Community Perceptions; Inappropriate Fire Regimes; Constrained Regenerative Capacity
Landholder Driven Planning for NRM	Medium	Land & Water Salinisation; Invasive Plants; Invasive Animals; Altered Hydrological Regimes; Soil Erosion; Inappropriate Water Use Practices; Recreational Pressures; Land Use Change; Direct Offsite Interactions; Misaligned Community Perceptions; Inappropriate Fire Regimes; Constrained Regenerative Capacity
Research to improve knowledge	High	Land & Water Salinisation; Invasive Plants; Invasive Animals; Altered Hydrological Regimes; Soil Erosion; Inappropriate Water Use Practices; Recreational Pressures; Land Use Change; Direct Offsite Interactions; Misaligned Community Perceptions; Inappropriate Fire Regimes; Constrained Regenerative Capacity
Asset condition monitoring and assessment	High	Land & Water Salinisation; Invasive Plants; Invasive Animals; Altered Hydrological Regimes; Soil Erosion; Inappropriate Water Use Practices; Recreational Pressures; Land Use Change; Direct Offsite Interactions; Misaligned Community Perceptions; Inappropriate Fire Regimes; Constrained Regenerative Capacity

Regional Delivery Partners

The individuals and organisations listed below have various roles to play within the Catchment Asset area in delivering and implementing the management actions described in previous sections that will in turn contribute to the aims and expected outcomes of the Mallee RCS. This list is not considered exhaustive and can be expected to change over time. More detail about the specific role and responsibilities of these entities with respect to the RCS can be found below.

- Aboriginal Affairs Victoria
- Aboriginal Communities (Traditional Owners & Organisations)
- Birdlife Mildura
- Catchment Management Authorities Mallee (VIC), Murray (NSW)
- Crown Frontage Licence Holders
- Department of Environment, Water and Natural Resources (SA)
- Department of Environment and Primary Industries (VIC)
- Department of Primary Industries & Regions SA (SA)
- Far West Anglers Association
- Lower Murray Water
- Mildura RSL Angling Club
- Mildura Rural City Council
- Millewa-Carwarp Landcare Group
- Murray Darling Basin Authority
- Murray Lower Darling Rivers Indigenous Nations
- NSW Department of Primary Industries Office of Water; Catchments & Lands; Fisheries
- NSW Office of Environment & Heritage
- NSW Transport & Maritime
- Parks Victoria
- Private Land Managers
- Regional Development Australia Loddon Mallee
- Research Bodies Arthur Rylah Institute, Invasive Animals CRC, Murray Darling Freshwater Research Centre
- South Australian Murray-Darling Basin Natural Resource Management Board
- Trust For Nature
- VicRoads
- VRFish
- Waterwatch
- Yelta Landcare Group

Community NRM Groups

This section provides a summary of these stakeholder groups that are active within this Catchment Asset. This is not an exhaustive list and will be updated on a regular basis. Further detail on the community NRM groups can be found in Attachment 1 – Mallee NRM Interest Groups.

Landcare Groups

The Landcare movement has been active in the Mallee since the late 1980s, with 23 active Landcare groups operating today. From our first group, Millewa-Carwarp (established 1989), to the newest, Cabarita (established 2012), Landcare has been instrumental in harnessing and promoting the interests of local communities in natural resource management.

Many groups initially formed due to the issues of rabbits and weeds. Salinity, soil conservation and biodiversity issues have evolved and continued to propel the Landcare movement into the 21st century. Our Landcare groups are keen to ensure that the legacy of Landcare continues and actively support a range of Junior Landcare Groups activities across the region.

Northern Mallee Landcare Consortium

The Northern Mallee Landcare Consortium represents the Millewa-Carwarp, Yelta, Cabarita, Red Cliffs and Kulkyne Way Landcare Groups

- Millewa-Carwarp
- Yelta

Non-aligned Landcare Groups

Cabarita

Other Groups

There are a broad range of community, industry, indigenous and specialist groups that have a role to play in NRM within this catchment asset. The tables in this section provide a listing of the groups currently known to be active within the asset area.

Traditional owner groups provide opportunities for our local indigenous communities to have input into how the regions landscapes are managed; and to enhance regional awareness of the cultural values inherent within these landscapes.

A wide range of special interest groups provide the means for individuals to become engaged in activities and programs which reflect their particular concerns. They also provide the region with a vital source of knowledge and understanding on specific issues.

The region's industry based groups have an important role in developing and promoting best practice for competitive and sustainable agricultural sectors.

Advisory groups to statutory bodies like CMA's and water authorities are important forums for both the community and the authorities involved. These groups provide management advice and community feedback to the convening authority while at the same time communicating with the wider community about the activities of the advisory group and the organisation in general.

Registered Aboriginal Parties

There is currently no registered aboriginal party determined for this Catchment Asset area

Special Interest and Recreational Groups

- BirdLife Mildura
- Murray Darling Association
- Murray Lower Darling Rivers Indigenous Nations
- Waterwatch

Industry Groups

- Victorian Farmers Federation (VFF)
- Victoria No-Till Farmers Association (VNTFA)
- Australian Table Grape Association Inc (ATGA)
- Dried Fruits Australia (DFA)
- Mallee Sustainable Farming Inc (MSF)
- Murray Valley Citrus Board (MVCB)
- Murray Valley Winegrowers' Inc (MVW)
- Pistachio Growers Association Incorporated (PGAI)
- Sunraysia Branch of the Victorian Apiarists Association

• Sunraysia Citrus Growers Inc (SCG)

Advisory Groups

- Aboriginal Reference Group (ARG)
- Mallee CMA Land and Water Advisory Committee (LWAC)
- Mallee CMA Technical Advisory Committees (TACs)
- Lower Murray Water Customer Service Advisory Committees (Rural)
- Goulburn-Murray Water Services Committees (WSCs)
- The Living Murray (TLM) Icon Site Community Reference Groups

Section 2 – Regional Assets in this Catchment Asset

Regional Assets in the Mallee NRM region were defined and applied in accordance with guidelines provided by DSE to support the Victorian Catchment Management Council RCS Guidelines. Each of the following sub-sections provides a background to these Regional Assets and the available information about their significance within this Catchment Asset. This follows on to an assessment of the relative value of the Regional Asset based on the criteria contained in Appendix 1. The Value of Regional Assets within a Catchment Asset

Rivers

The Murray River is Australia's largest river and firmly entrenched in the national psyche. It forms the northern boundary of the Victorian Mallee NRM region, and the border between Victoria and NSW. While management of the main river is the statutory responsibility of NSW, Victoria is responsible for the management of its southern floodplain from the 1881 winter level mark. The River's floodplain, anabranches and associated wetland systems dominate the area.

Apart from the reaches of Murray River itself between Merbein and the South Australian Border, there are two Murray anabranch systems within this Catchment Asset. They are the Lindsay River (one of the largest Murray anabranches systems in Victoria) and Wallpolla Creek. Within each of these systems are a number of smaller, interlinking waterways that wind across their respective floodplains. In total, this Catchment Asset contains over 770 kilometres of waterways

The Living Murray Initiative is a program of measures established by the Murray-Darling Basin Ministerial Council with a focus on achieving environmental benefits for six significant ecological assets through the recovery of up to 500 gigalitres over five years. The Murray River channel is one of the six Icon sites included in the Living Murray Initiative.

The Mallee Waterway Strategy (2014-22) identifies priority reaches for future management using an Asset Based Approach; facilitating targeted planning and implementation processes to deliver greatest social, cultural and economic returns on our efforts. Each River asset occurring within this Catchment Asset is ascribed a priority of 'high', 'medium', 'low', or 'additional' under this framework.

Based on the available data concerning their extent, quality and policy significance within this landscape, the Rivers within this Catchment Asset are considered to be of **very high** value.




Figure 1: River reaches

Wetlands

This Catchment Asset encompasses the wetland systems of Lindsay and Wallpolla Islands and Lake Wallawalla. Together they occupy almost 25,000 hectares of the Catchment Asset area and they all lie within the boundaries of the Murray Sunset National Park. There is a further 15,000 hectares of wetlands and floodplain that lay outside these systems yet within the Catchment Asset.

The Lindsay and Wallpolla Islands and Lake Wallawalla systems are each nationally recognised in the Directory of Important Wetlands and named in JAMBA and CAMBA as habitat for international migratory birds. They also form the upstream component of the Chowilla Floodplain Complex which straddles the Victorian and South Australian Borders and is one of the six Living Murray Icon Sites.

The Mallee Waterway Strategy (2014-22) identifies priority wetlands for future management using an Asset Based Approach; facilitating targeted planning and implementation processes to deliver greatest social, cultural and economic returns on our efforts. Each Wetland asset occurring within this Catchment Asset is ascribed a priority of 'high', 'medium', 'low', or 'additional' under this framework.

Based on the available data concerning their extent, quality and policy significance within this landscape, the Wetlands within this Catchment Asset are considered to be of **very high** value.




Figure 2: Wetlands

Threatened Species and Communities

A significant range of flora and fauna species that are listed in Federal and State instruments have been observed and recorded within the area of this catchment asset over time. These species are listed in the tables that follow along with their current status as described in the listing instrument. The listing instruments are: Environmental Protection and Biodiversity Conservation (EPBC) Act 1999; Flora and Fauna Guarantee (FFG) Act 1998 and the Victorian Threatened Species Advisory Lists issued by the Department of Environment and Primary Industries (DEPI) (Flora: 2005, Fauna: 2013). The species names included in the following tables (Table 4 &

Table 5) may not reflect the full range of threatened flora and fauna species that may inhabit the catchment area. Rather they reflect only species whose presence has been observed and recorded by suitably qualified observers since 1980.

This Catchment Asset is one of a number in the Mallee Region that contain remnant areas of the nationally listed ecological community, 'Buloke Woodlands of the Riverina and Murray-Darling Depression Bioregions'. This ecological community is listed as 'endangered' under the Environment Protection and Biodiversity Conservation Act 1999 (EPBC Act).

Lindsay Island supports scattered stands of the 'semi-arid woodland' and 'semi-arid chenopod woodland' ecological vegetation classes that include within them stands of 'Buloke Woodland', a vegetation community listed as 'endangered' under the national EPBC Act 1999. These stands are an unusual variant of those found elsewhere in the Mallee because the depleted White Cypress Pine (Callitris glaucophylla) replaces Slender Cypress Pine (Callitris gracilis) as one of the dominant cospecies.

The river reaches, anabranches and floodplain wetlands contained within this Catchment Asset area provide habitat for some of the members of the Lowland Riverine Fish Community of the Southern Murray–Darling Basin (see

Table 6). This is a community of fish species that is listed under the Victorian FFG Act and so qualifies as a threatened ecological community in this context. The Mallee Bird Community is another listed ecological community under the FFG Act. However, only one species (Gilbert's Whistler) from this group has been recorded within the catchment since 1980 and this species is not considered to be threatened.

Based on the available data concerning their extent, quality and policy significance within this landscape, the Threatened Species and Communities within this Catchment Asset are considered to be of high value.

Table 4: Threatened Flora Species observed since 1980

Scientific Name	Common Name	EPBC	FFG	DSE Advisory List
Abutilon fraseri	Dwarf Lantern-flower		Listed	Endangered
Alternanthera nodiflora	Common Joyweed			Poorly Known
Amaranthus macrocarpus var. macrocarpus	Dwarf Amaranth			Vulnerable
Angianthus brachypappus	Spreading Angianthus			Vulnerable
Asperula gemella	Twin-leaf Bedstraw			Rare
Atriplex acutibractea	Pointed Saltbush			Rare
Atriplex acutibractea subsp. karoniensis	Pointed Saltbush			Rare
Atriplex limbata	Spreading Saltbush		Listed	Vulnerable
Atriplex lindleyi subsp. conduplicata	Baldoo			Rare
Atriplex nummularia subsp. omissa	Dwarf Old-man Saltbush			Rare
Atriplex pseudocampanulata	Mealy Saltbush			Rare
Atriplex rhagodioides	Silver Saltbush		Listed	Vulnerable
Atriplex vesicaria subsp. macrocystidia	Bladder Saltbush			Poorly Known
Atriplex vesicaria subsp. minor	Bladder Saltbush			Poorly Known
Austrobryonia micrantha	Mallee Cucumber			Rare
Austrostipa pilata	Prickly Spear-grass			Vulnerable
Bergia trimera	Small Water-fire			Vulnerable
Bromus arenarius	Sand Brome			Rare
Calandrinia volubilis	Twining Purslane			Rare
Calostemma purpureum s.l.	Garland Lily			Rare
Calotis cuneifolia	Blue Burr-daisy			Rare
Cardamine moirensis	Riverina Bitter-cress			Rare
Centipeda nidiformis	Cotton Sneezeweed			Rare
Centipeda pleiocephala	Tall Sneezeweed			Endangered
Centipeda thespidioides s.l.	Desert Sneezeweed			Rare

Convolvulus clementii	Desert Bindweed		Vulnerable
Craspedia haplorrhiza	Plains Billy-buttons		Poorly Known
Crinum flaccidum	Darling Lily	Listed	Vulnerable
Cullen cinereum	Hoary Scurf-pea	Listed	Endangered
Cynodon dactylon var. pulchellus	Native Couch		Poorly Known
Cyperus pygmaeus	Dwarf Flat-sedge		Vulnerable
Cyperus victoriensis	Yelka		Poorly Known
Dissocarpus biflorus var. biflorus	Twin-flower Saltbush		Rare
Duma horrida subsp. horrida	Spiny Lignum		Rare
Elachanthus glaber	Smooth Elachanth		Rare
Eleocharis pallens	Pale Spike-sedge		Poorly Known
Epaltes cunninghamii	Tall Nut-heads		Vulnerable
Eragrostis australasica	Cane Grass		Vulnerable
Eragrostis lacunaria	Purple Love-grass		Vulnerable
Eragrostis setifolia	Bristly Love-grass		Vulnerable
Eremophila bignoniiflora	Bignonia Emu-bush	Listed	Vulnerable
Eremophila divaricata subsp. divaricata	Spreading Emu-bush		Rare
Eremophila maculata subsp. maculata	Spotted Emu-bush		Rare
Eremophila polyclada	Twiggy Emu-bush		Vulnerable
Eremophila sturtii	Narrow-leaf Emu-bush	Listed	Endangered
Euphorbia tannensis subsp. eremophila	Desert Spurge	Listed	Endangered
Fimbristylis velata	Veiled Fringe-sedge		Rare
Isolepis australiensis	Inland Club-sedge		Poorly Known
Lepidium fasciculatum	Bundled Peppercress		Poorly Known
Lepidium papillosum	Warty Peppercress		Poorly Known
Lepidium pseudohyssopifolium	Native Peppercress		Poorly Known
Leptochloa fusca subsp. fusca	Brown Beetle-grass		Rare
Maireana aphylla	Leafless Bluebush		Poorly Known
Maireana sedifolia	Pearl Bluebush		Rare
Malacocera tricornis	Goat Head		Rare
Minuria cunninghamii	Bush Minuria		Rare
Minuria denticulata	Woolly Minuria		Rare
Minuria integerrima	Smooth Minuria		Rare

Neobassia proceriflora	Soda Bush		Endangered
Nicotiana goodspeedii	Small-flower Tobacco		Rare
Phyllanthus lacunarius	Lagoon Spurge		Vulnerable
Picris squarrosa	Squat Picris		Rare
Rhodanthe polygalifolia	Milkwort Sunray		Rare
Rorippa eustylis	Dwarf Bitter-cress		Rare
Rumex crystallinus s.s.	Glistening Dock		Vulnerable
Sclerolaena decurrens	_		Vulnerable
	Green Copperburr		
Sclerolaena divaricata	Tangled Copperburr		Poorly Known
Sclerolaena intricata	Poverty Bush		Vulnerable
Sclerolaena muricata var. muricata	Black Roly-poly		Poorly Known
Senecio cunninghamii var. cunninghamii	Branching Groundsel		Rare
Senecio platylepis	Toothed Groundsel		Rare
Sida intricata	Twiggy Sida		Vulnerable
Solanum lacunarium	Lagoon Nightshade		Vulnerable
Swainsona greyana	Hairy Darling-pea	Listed	Endangered
Swainsona microphylla	Small-leaf Swainson-pea		Rare
Swainsona reticulata	Kneed Swainson-pea	Listed	Vulnerable
Tecticornia triandra	Desert Glasswort		Rare
Templetonia egena	Round Templetonia		Vulnerable
Tetragonia eremaea s.s.	Desert Spinach		Poorly Known
Tetragonia moorei	Annual Spinach		Poorly Known
Velleia arguta	Grassland Velleia		Rare
Vittadinia condyloides	Club-hair New Holland Daisy		Rare
Wahlenbergia tumidifructa	Mallee Annual-bluebell		Rare
Zygophyllum simile	White Twin-leaf		Rare

Table 5: Threatened Fauna Species observed since 1980

Scientific Name	Common Name	EPBC	FFG	DSE Advisory List
Alcedo azurea	Azure Kingfisher			Near Threatened
Anas rhynchotis	Australasian Shoveler			Vulnerable
Ardea intermedia	Intermediate Egret		Listed	Endangered

Ardea modesta	Eastern Great Egret L		Listed	Vulnerable
Aythya australis	Hardhead			Vulnerable
Biziura lobata	Musk Duck			Vulnerable
Burhinus grallarius	Bush Stone-curlew Listed		Listed	Endangered
Charadrius australis	Inland Dotterel			Vulnerable
Chelodina expansa	Broad-shelled Turtle		Listed	Endangered
Chrysococcyx osculans	Black-eared Cuckoo			Near Threatened
Climacteris affinis	White-browed Treecreeper		Listed	Vulnerable
Climacteris picumnus victoriae	Brown Treecreeper (southeastern ssp.)			Near Threatened
Denisonia devisi	De Vis' Banded Snake			Critically Endangered
Dromaius novaehollandiae	Emu			Near Threatened
Egretta garzetta nigripes	Little Egret		Listed	Endangered
Falco hypoleucos	Grey Falcon		Listed	Endangered
Falco subniger	Black Falcon	Black Falcon		Vulnerable
Furina diadema	Red-naped Snake		Listed	Vulnerable
Geopelia cuneata	Diamond Dove		Listed	Near Threatened
Haliaeetus leucogaster	White-bellied Sea-Eagle	l Sea-Eagle Lis		Vulnerable
Hydroprogne caspia	Caspian Tern	Caspian Tern Lis		Near Threatened
Litoria raniformis	Growling Grass Frog	g Grass Frog Vulnerable		Endangered
Lophocroa leadbeateri	Major Mitchell's Cockatoo	00		Vulnerable
Maccullochella peelii	Murray Cod	Vulnerable	Listed	Vulnerable
Macquaria ambigua	Golden Perch			Near Threatened
Melanodryas cucullata cucullata	Hooded Robin		Listed	Near Threatened
Melanotaenia fluviatilis	Murray-Darling Rainbowfish		Listed	Vulnerable
Morelia spilota metcalfei	Carpet Python		Listed	Endangered
Ninox connivens connivens	Barking Owl		Listed	Endangered
Nycticorax caledonicus hillii	Nankeen Night Heron	Night Heron		Near Threatened
Oreoica gutturalis gutturalis	Crested Bellbird	Crested Bellbird Listed		Near Threatened
Oxyura australis	Blue-billed Duck	e-billed Duck List		Endangered
Parasuta spectabilis	Port Lincoln Snake	ort Lincoln Snake Lis		Vulnerable
Phalacrocorax varius	Pied Cormorant			Near Threatened
Planigale gilesi	Gile's Planigale		Listed	Near Threatened
Platalea regia	Royal Spoonbill			Near Threatened

Polytelis anthopeplus monarchoides	Regent Parrot Vulnerable		Listed	Vulnerable
Porzana pusilla palustris	Baillon's Crake		Listed	Vulnerable
Pseudonaja aspidorhyncha	Patch-nosed Brown Snake			Near Threatened
Ptilonorhynchus maculatus	Spotted Bowerbird		Listed	Critically Endangered
Pyrrholaemus brunneus	Redthroat		Listed	Endangered
Rhynchoedura ornata	Beaked Gecko		Listed	Critically Endangered
Sminthopsis crassicaudata	Fat-tailed Dunnart			Near Threatened
Stictonetta naevosa	Freckled Duck		Listed	Endangered
Stiltia isabella	Australian Pratincole			Near Threatened
Struthidea cinerea	Apostlebird		Listed	
Tandanus tandanus	Freshwater Catfish		Listed	Endangered
Todiramphus pyrropygia pyrropygia	Red-backed Kingfisher			Near Threatened
Tringa nebularia	Common Greenshank			Vulnerable
Turnix pyrrhothorax	Red-chested Button-quail		Listed	Vulnerable
Varanus varius	Lace Monitor			Endangered

Table 6: Lowland Riverine Fish Community of the Southern Murray–Darling Basin as listed under FFG Act (including individual species listing and number of records since 1980 from Victorian Biodiversity Atlas)

Scientific Name	Common Name	Records since 1980	EPBC	FFG	DSE Advisory List
Ambassis agassizii	Agassiz's Chanda Perch	No Records		Listed	Regionally Extinct
Bidyanus bidyanus	Silver Perch	No Records		Listed	Vulnerable
Craterocephalus fluviatilis	Murray Hardyhead	No Records	Endangered	Listed	Critically Endangered
Craterocephalus stercusmuscarum fulvus	Non-specked Hardyhead	1			
Galaxias rostratus	Flat-headed Galaxias	No Records			Vulnerable
Hypseleotris klunzingeri	Western Carp Gudgeon	11			
Maccullochella macquariensis	Trout Cod	No Records	Endangered	Listed	Critically Endangered
Maccullochella peelii peelii	Murray Cod	4	Vulnerable	Listed	Vulnerable
Macquaria ambigua	Golden Perch	33			Near Threatened
Macquaria australasica	Macquarie Perch	No Records	Endangered	Listed	Endangered
Melanotaenia fluviatilis	Murray-Darling Rainbowfish	5		Listed	Vulnerable
Mogurnda adspersa	Southern Purple- spotted Gudgeon	No Records			Regionally Extinct
Nematoalosa erebi	Bony Bream	22			
Philypnodon grandiceps	Flat-headed Gudgeon	1			
Tandanus tandanus	Freshwater Catfish	1		Listed	Endangered

Terrestrial Habitat

A little over ninety seven percent of this Catchment Asset area is covered by native vegetation. The area is also amongst the more diverse in terms of Ecological Vegetation Classes (EVC) with almost half of the Mallee's EVCs being found within the asset. The catchment asset lies almost entirely within the Victorian portion of the Interim Biogeographic Regionalisation for Australia (IBRA) sub-region known as the Murray Scroll Belt (seeFigure 3). This sub-region is itself part of the Riverina IBRA bioregion which extends eastwards upriver to take in the lower Murrumbidgee, mid Murray and lower Goulburn & Broken River valleys. However, the catchment asset also takes in a small section of the adjacent Murray Mallee IBRA sub-region which is part of the Murray-Darling Depression IBRA bioregion that takes in the lower south western portion of the Murray-Darling Basin, spread across Victoria, South Australia and New South Wales.

Vegetation on frontages adjacent to the River is comprised predominantly of River Red Gum (Eucalyptus camaldulensis) communities with the floodplains further back characterised by Black Box (Eucalyptus largiflorens) woodlands, lignum shrublands and grasslands. The floodplain area adjoins large, open areas of semi-arid grasslands and chenopod shrublands.

The table below lists each of the Catchment Asset's EVCs along with the IBRA subregion within which they are contained, their Bioregional Conservation Status (BCS) and their area in hectares.

Table 7: Area of Ecological Vegetation Classes and their Bioregional Conservation Status

Ecological Vegetation Class	IBRA Subregion	Conservation Status	Area (hectares)
Alluvial Plains Semi-arid Grassland	Murray Scroll Belt	Vulnerable	3368.90
Bare Rock/Ground	Murray Mallee	Not Applicable	1.93
	Murray Scroll Belt	Not Applicable	477.01
Chenopod Mallee	Murray Mallee	Vulnerable	431.17
	Murray Scroll Belt	Vulnerable	79.39
Disused Floodway Shrubby Herbland	Murray Scroll Belt	Endangered	22.81
Floodplain Grassy Wetland	Murray Scroll Belt	Endangered	20.68
Floodway Pond Herbland	Murray Scroll Belt	Depleted	319.17
Grassy Riverine Forest	Murray Scroll Belt	Depleted	898.42
Grassy Riverine Forest/Floodway Pond Herbland Complex	Murray Scroll Belt	Depleted	283.88
Intermittent Swampy Woodland	Murray Scroll Belt	Depleted	3384.37
Lake Bed Herbland	Murray Scroll Belt	Vulnerable	219.46
Lignum Shrubland	Murray Scroll Belt	Least Concern	10719.40
Lignum Swamp	Murray Scroll Belt	Vulnerable	1025.36
Lignum Swampy Woodland	Murray Scroll Belt	Depleted	4470.83
Low Chenopod Shrubland	Murray Scroll Belt	Depleted	10126.14
Riverine Chenopod Woodland	Murray Mallee	Depleted	25.14
	Murray Scroll Belt	Depleted	15657.59
Semi-arid Chenopod Woodland	Murray Scroll Belt	Depleted	4148.16
	Murray Mallee	Vulnerable	139.74
Semi-arid Woodland	Murray Mallee	Vulnerable	34.86
	Murray Scroll Belt	Vulnerable	122.85
Shallow Freshwater Marsh	Murray Scroll Belt	Vulnerable	68.78
Shrubby Riverine Woodland	Murray Mallee	Least Concern	7.86
	Murray Scroll Belt	Least Concern	4345.73
Spike-sedge Wetland	Murray Scroll Belt	Vulnerable	55.80
Sub-saline Depression Shrubland	Murray Scroll Belt	Vulnerable	872.99

Water Body - Fresh	Murray Scroll Belt	Not Applicable	583.62
Woorinen Sands Mallee	Murray Scroll Belt	Depleted	0.28




Figure 3: Bioregional conservation status of EVCs and IBRA sub-regions

DEPI's NaturePrint model is a landscape scale spatial planning mechanism that provides an analysis of biodiversity values across Victoria and combines the available information about biodiversity values, threatening processes and ecosystem function. It provides an analysis of the ecological value of a portion of landscape compared across the entire Victorian landscape. It is based on the following measures:

- Mathematical models of species distributions and habitats;
- The condition of these habitats:
- · Pathways for connectivity across landscapes;
- Connectivity potential and recoverability;
- Threats to species persistence.

For background information about the model, its output and its significance, refer to: http://www.depi.vic.gov.au/environment-and-wildlife/biodiversity/natureprint.

The NaturePrint model considers that over 91% of this Catchment Asset contains terrestrial habitat that is of the highest ecological value to Victoria and therefore high priority for protection and conservation.

Based on the available data concerning their extent, quality and policy significance within this landscape, the Terrestrial Habitat within this Catchment Asset is considered to be of **very high** value.




Figure 4: Terrestrial Habitat given high ecological value by NaturePrint Version 3

Soils

The soils of this asset area are predominantly the heavy grey cracking floodplain clays that form a grouping known as Vertosols. These are coarsely structured soils, thought to be derived from ancient lake deposits, that typically sodic, gypseous and somewhat saline and therefore not typically suited to agriculture. Further back from the river country, on the higher ground, are the texture contrast soils known as Sodosols. These are typically red clays and earths, often somewhat saline, which are often vulnerable to erosion. Despite this, only a tiny proportion (1.6% or 776 hectares) of soils within the asset area has been identified as being of high susceptibility to wind erosion.

A recent survey of the Murray River frontage in this asset area identified a number of locations where the river bank soils were subject to a number of different forms of water erosion. Whilst many could be considered as normal fluvial responses to water flows, some may have been exacerbated by external processes such as high river flows and reduced vegetation cover.

The soils of this Catchment Asset area are predominantly valued for their capacity to provide for terrestrial habitat and the support the production of food and fibre. However, there is also an understanding that the structure of the soil and its natural capacity to resist threatening processes such as wind erosion is also of substantial value.

There is currently no accepted framework to categorise our soil asset in terms of value from lower to higher based purely upon the social, economic and environmental values and services they provide. Such a framework has only been considered to date in terms of the threatening processes that may impinge on the capacity of our soils asset to provide the services we have come to expect. This will be discussed later in this document. Despite this limitation, the most appropriate course of action is to consider the relative value of the soil asset according to its intrinsic capacity to support the other regional assets within this catchment asset as the basis for applying a value category.

Therefore, the Soils within this Catchment Asset are considered to be of high value.




Figure 5: Dominant Soil Types

Agricultural Land

Only 10% of the catchment asset is given over to agricultural activities. Irrigated agriculture is confined largely to the area around Lindsay Point while a large area of mixed dryland farming is pursued to the east of Lake Cullulleraine.

The agricultural land of this Catchment Asset area is mainly valued for its capacity to support the production of food and fibre. This in turn provides for a range of other social and economic services that are important to the Mallee region. In addition, when these services are available, they support the local community in its efforts to maintain and enhance not only their agricultural land but other regional assets. There is currently no accepted framework to categorise our agricultural land asset in terms of value from lower to higher based purely upon the social and economic values and services they provide. Such a framework has only been considered to date in terms of the threatening processes that may impinge on the capacity of our agricultural land asset to provide the services we have come to expect. This will be discussed later in this document. As a result, the most appropriate course of action is to consider the relative value of the agricultural land asset at a regional scale rather than at a catchment asset scale and in accordance to its capacity to provide and support social and economic values.

Therefore, the Agricultural Land within this Catchment Asset and across the region is considered to be of high value.




Figure 6: Agricultural land use

Groundwater

Groundwater aquifers that lie underneath this Catchment Asset area are not known to be utilised to provide a water resource for human uses. Therefore, the Mallee's groundwater asset is not of significance to this Catchment Asset.

Cultural Heritage

Most of the floodplain has high cultural heritage for local indigenous communities, particularly areas of river frontage and wetlands. Throughout this entire area the large concentration of burials is very unique to Victoria. There are also a significantly large amount of meeting places, ceremonial places, scarred trees, and artefact sites. These sites show that Aboriginal people inhabited the area for tens of thousands of years. Over 150 Indigenous Heritage Sites in the Lindsay-Wallpolla Island area have been listed with Aboriginal Affairs Victoria, though it is understood that more exist.

Cultural heritage sites in this Catchment Asset area are predominantly valued for the connection they provide to the community between the social and cultural landscape that we currently enjoy and the stories, ideas and history of how that came to be. These sites provide individuals and families with a physical representation of their connection to the Mallee landscape and their place in it. These connections are extremely important to these people and, in turn, can reflect upon how they value and relate to the Mallee landscape that they are familiar with today. Therefore it is important to all that these sites are managed and protected in a manner that is sympathetic to those connections and values. Heritage significance is typically assessed under a three-tier legislative system that determines the local, state, national or commonwealth significance of an item. Heritage items can also be included on non-statutory listings (such as the National Trust classifications). However, there is no single and simple approach to categorise our cultural heritage asset in terms of value to the region. Tradition distinctions between different types of value (such as economic, social and environmental values) can be determined but this data is only patchily available given the number and range of cultural heritage sites across the Mallee. The most appropriate course of action at this point is to give equal value to all recognised cultural heritage sites in the Mallee regardless of their age, form and historical background.

Therefore, Cultural Heritage within this Catchment Asset and across the region is considered to be of high value. Any further processes undertaken to prioritise and target management actions will rely on assessments of risk likelihood and impact to those sites.




Figure 7: Areas of Cultural Value

Community Capacity

Community capacity is an important asset as positive and long-lasting NRM outcomes are dependent on an active, willing and capable community. In order to achieve regional scale NRM outcomes, people have to play a major role, often in partnership with government and industry, in the ongoing conservation and sustainable use of our Regional Assets This requires Mallee communities to have the capacity to work cooperatively, apply economic resources, use networks and acquire and use relevant knowledge and information to achieve NRM outcomes. Maintaining and improving this capacity is dependent not only on the financial, physical and natural resources contained within or available to a community but also its social resources.

Community capacity therefore requires ongoing development, conservation and management as its relative condition can influence the achievement of all other biophysical goals and actions outlined within the RCS.

Our communities' capacity is defined by their characteristics and resources which, when combined, determine their ability to identify, evaluate and address key issues.

Characteristics of strong regional community capacity include:

- Individuals within the community being aware of regional NRM issues, and understanding the link between these issues and the long-term viability of the community
- Natural resource managers and users being able and willing to access the necessary information, data and science – biophysical, social and economic – to make sound NRM decisions
- Natural resource managers and users being equipped with or having access to, the necessary technical, people management, project management and planning skills to Social, participate

- in the development and implementation of sustainable NRM at the property, local and regional scales
- Community being engaged and motivated, and exercising ownership over NRM decision making processes and effectively implementing actions arising from these processes (DAFF 2006).

This catchment asset area is very sparsely populated with no distinct centres of human occupation, rather just scattered homes that support farming enterprises and similar activities. However, this area is heavily used as a recreation resource and tourism destination by populations both within and external to the Mallee. Therefore, while the resident community is of critical importance to the ongoing management of the catchment asset, it is this wider population of occasional users whose capacity for effective and sustainable natural resource management is potentially quite influential in the long term by virtue of its greater size.

This catchment asset area is well serviced by community driven NRM organisations which reflects well on the capacity of the local community to contribute to effective NRM outcomes. There are two active Landcare groups (Millewa-Carwarp; and Yelta) whose areas of influence lie partially or wholly within the asset area.

The asset area is one of strongholds of Mallee Sustainable Farming, a highly respected not-for-profit agricultural and research and extension organisation that is led by and works for cropping farmers in the northern half of the Mallee, south-western NSW and the Riverland of SA.

Trust for Nature is another important contributor to NRM within this asset area. This not for profit organisation is a Victorian Government statutory body that fosters environmental conservation principles through the use of conservation covenants. One of its functions is the acquisition of land either temporarily or permanently in order to apply perpetual conservation covenants to the land to benefit biodiversity. Trust for Nature owns the 30,000 hectare Ned's Corner Station and much of this property lies within this Catchment Asset area.

A complete listing of all the known community NRM organisations within the catchment asset can be found in the Community NRM Groups section.

Whilst there are a number of methods available to evaluate the relative 'condition' of our community's capacity for NRM there is currently no accepted framework to categorise the community capacity asset in terms of value from low to high based upon the values and services it may provide. As a result, the most appropriate course of action is to consider the relative value of community capacity for NRM in terms of its potential contribution to NRM outcomes at a regional scale and what those NRM outcomes might be (and how sizeable the resources required to achieve them) if that contribution was not forthcoming. Under those circumstances, it is relatively easy to come to the conclusion that, without the communities' investment into NRM through exercising that community capacity asset, the scale and success of our NRM outcomes will be curtailed while their cost in resources increases. Such a situation would not benefit the Mallee region or its Regional Assets.

Therefore, Community Capacity within this Catchment Asset and across the region is considered to be of high value.




Figure 8: State Suburbs

Section 3 – Threatening Processes in this Catchment Asset

Threatening processes in the Mallee NRM region were defined and applied as part of the development of Mallee Regional Catchment Strategy by the Regional Catchment Strategy Steering Committee. Each of the following sub-sections provides a background to these threatening processes, available information about their form within this Catchment Asset and some discussion regarding the impact of these processes on our regional assets. This follows on to an assessment of the relative threat level posed by each the threatening processes based on the criteria contained in Appendix 2. *Categorising Threatening Processes*.

Land & water salinisation

Land and water salinisation is defined as: "Salinisation of the soil and water resources through the displacement of salt as a result of natural and human induced drivers."

Salinisation of soils is caused by discharge of groundwater to the atmosphere by direct transpiration of groundwater by vegetation or by capillary rise from the water table through the soil profile to either the soil surface (evaporation) or the bottom of the plant root zone. Salt contained in the water that is evaporated and/or transpired is left behind and accumulates in the soil. High rates of groundwater discharge, and hence salt accumulation, only occur in areas where water tables are shallow (typically less the 10 m, with the most severe effects when the water table is less than 5 m).

There are many naturally saline wetlands and wetland complexes in the Mallee that are situated in areas where the underlying water tables are naturally shallow. Better known examples of these natural systems include the Tyrrell Basin, Raak Plain and the Pink Lakes. These are natural saline systems that have historically developed and been maintained in a balance that provided for stable ecological communities. However, agricultural development in the last century has altered the long term balance of these systems. The removal of much of the Mallee vegetation and the use of land management techniques that were comparatively inefficient users of rainfall (compared to Mallee woodlands) allowed deep drainage past the root zone. Depending on the location, this deep drainage either manifest itself as localised discharge zones in dune swales or contributed to further shallowing of the regional water table. As water tables came closer to the surface, more wetlands and wetland ecosystems were threatened by increasing salinity and increasing areas of salinised landscape.

The incremental improvement in agricultural land management techniques to maximise water use efficiency and minimise through drainage over the last two decades coupled with the broad scale effect of the so-called Millennium Drought have begun to swing the balance back towards a more neutral status. Water table depths have fallen and correspondingly so have salinity impacts in the landscape.

Despite these recent advances, the risks remain. Many of the triggers in the landscape that created the salinisation problem still remain. Most notable of these are the altered flow regimes in our watercourses and wetlands and the large, open areas cleared of perennial Mallee vegetation. These are historical landscape modifications that are not expected to be rectified since they have provided for the bulk of the Mallee's resultant economic and social activity. Lapses in our current regimes of effective land and water management coupled with a changing climate can easily contribute to the return of a significant period of land and water salinisation.

This catchment asset includes a large area of naturally saline landscape situated to the west of Lake Cullulleraine which accounts for a little over 10% of the total catchment asset area that has been identified as having induced saline surfaces. Classification of induced and natural saline surfaces in 2009 indicates there is only 99 hectares of induced saline surfaces in the catchment asset area (Mock & Grinter, 2009). This induced salinity is generally confined to a small number of wetlands in the eastern end of the asset area. These areas of induced saline surfaces represent only a 1.2% increase in the area of saline surfaces over and above that which would be considered 'natural' (see Figure 9).

Depth to the regional water table and the general trends in groundwater levels were modelled in 2010 throughout the Mallee based on analysis of available groundwater data. A little over 3% of the catchment asset area has typically saline groundwater within 2 metres of the ground surface while more

than 60% has saline groundwater within 5 metres of the ground surface. Therefore, over 30,000 hectares of the catchment asset area is considered to have a moderate or higher risk of salinisation (see Figure 10).

Given the current status of land and water salinisation in the Catchment asset area and consideration of the future potential of this threatening process to produce negative impacts within the area, the priority of action with respect to each Regional Asset has been determined. They are listed in the table below along with the summary of the logic behind the priority determination.

Table 8: Prospective Priority of Action to address Land and Water Salinisation upon Regional Assets

Regional Asset	Priority of Action	Summary Logic
Rivers	HIGH	No aquatic or riparian weeds identified to date; no rivers in high priority IPAWS asset area
Wetlands	HIGH	No aquatic or riparian weeds identified to date; some priority wetlands within high priority IPAWS asset area
Threatened Species and Communities	MEDIUM	Buloke woodlands are an endangered ecological community and lie within the high priority IPAWS asset area
Terrestrial Habitat	MEDIUM	Most remnant habitat in asset area is small and linear in shape; much lies within the high priority IPAWS asset area
Soils	HIGH	Most identified weeds in the asset area do not pose a significant threat to soil health at this time
Agricultural Land	MEDIUM	Most identified weeds in the asset area only represent localised threat to agriculture, primarily of nuisance value
Groundwater	NONE	This regional asset is not present within this catchment asset
Cultural Heritage	NONE	Most identified weeds in the asset area do not pose a significant threat to cultural heritage sites at this time
Community Capacity	NONE	Invasive plants do not pose a significant threat to community capacity at this time




Figure 9: Natural and Induce Saline Surfaces




Figure 10: Estimated Depth to the Water table

Invasive Plants

Currently there are 25 State Prohibited Weeds, 4 Regionally Prohibited Weeds and 14 Regionally Controlled Weeds listed under the Catchment and Land Protection (CaLP) Act for the Mallee region. In addition to the conventional ways that weeds spread (e.g. wind and water) in the Mallee, disturbance from pest animals such as rabbits, goats and pigs also create good conditions for weed growth and contribute to weed spread.

Invasive plants in the Mallee threaten both biodiversity and the productive capacity of land. In many cases, weed infestation has changed the composition and character of the ecological landscape. Weed infestations also impact on the use of popular recreational spots in riparian areas, affecting aesthetic values and limiting access. Environmental weeds compete with established and regenerating indigenous species, threatening the quality and extent of native vegetation and reducing the diversity and availability of habitat. Environmental weeds present a significant risk to remnant roadside vegetation, as the 'edge effect' is particularly difficult to manage in these long, thin areas.

Agricultural weeds cause significant losses for horticulture and dryland farming through competition with crops and by reducing the quality of produce. Their control increases the cost of production and, in extreme cases, can diminish the productive capability of the land.

While the presence of recognised weed species in the asset area is almost certain, there is insufficient data available to adequately describe the current scope and scale of pest plant populations for the purposes of effective regional control. The available spatial data on invasive plant infestations dates back to 2007 and was generated as part of the Priority Weed Mapping project in 2008. This information is obviously now far out of date. The current information and response capacity relies heavily on local activity at a local scale in response to local weed infestations. This approach, while often very successful, tends to preclude centralised management and control at a regional scale. Therefore, the priority of action to address invasive plants is strongly influenced by local scale data derived from local planning instruments and informal community consultation at the local level. On the other hand, limited knowledge and understanding of the vulnerability of the Mallee landscape to future weed incursions is available given the current listing of known invasive plants, the current understanding of their biological requirements and the planning and response framework provided by regional scale plans such as the Mallee Invasive Plants and Animals Management Strategy (IPAWS).

The IPAWS identifies 21 regional assets (based on geographic areas) across the Mallee region. This Catchment Asset area contains four of these geographic assets: Agricultural Land; Tyrrell Creek and Lake Tyrrell; Lalbert Creek and Lake Timboram; and the Southern Buloke Woodland Community. Of these, only the Southern Buloke Woodland Community was considered to be a high priority for onground action to manage invasive plants and animals. The remaining three assets were identified as areas where monitoring for future action and maintaining previous management gains was considered sufficient.

Using this combination of information gathering approaches and the application of a complimentary consultative process, the priority of action for dealing with invasive plants with respect to each Regional Asset has been determined. They are listed in the table below along with the summary of the logic behind the priority determination.

Table 9: Prospective Priority of Action to address Invasive Plants upon Regional Assets

Regional Asset	Priority of Action	Summary Logic
Rivers	HIGH	No aquatic or riparian weeds identified to date; no rivers in high priority IPAWS asset area
Wetlands	HIGH	No aquatic or riparian weeds identified to date; some priority wetlands within high priority IPAWS asset area
Threatened Species and Communities	HIGH	Buloke woodlands are an endangered ecological community and lie within the high priority IPAWS asset area
Terrestrial Habitat	HIGH	Most remnant habitat in asset area is small and linear in shape; much lies within the high priority IPAWS asset area
Soils	HIGH	Most identified weeds in the asset area do not pose a significant threat to soil health at this time
Agricultural Land	HIGH	Most identified weeds in the asset area only represent localised threat to agriculture, primarily of nuisance value
Groundwater	NONE	This regional asset is not present within this catchment asset
Cultural Heritage	HIGH	Most identified weeds in the asset area do not pose a significant threat to cultural heritage sites at this time
Community Capacity	NONE	Invasive plants do not pose a significant threat to community capacity at this time




Figure 11: Priority Weed Infestations Recorded Between 2005 and 2007




Figure 12: Regional Assets from the Mallee Invasive Plants and Animal Strategy

Invasive Animals

Invasive animals in terms of the Mallee RCS are regarded as those established invasive pest animals as defined in Part 8 of the CaLP Act 1994

Australia is host to 56 invasive vertebrate animal species. Of these, the ones with the most impact (in order of damage estimates) are: European Red Fox, feral cats, rabbits, feral pigs, wild dogs, the house mouse, carp and goats. The Mallee region hosts populations of all of these species. However, of most significance to the Mallee is the European Red Fox and Rabbits – both are declared under the Catchment and Land Protection (CaLP) Act 1994.

Invasive species are considered to be pests when they have, or have the potential to have, an undesirable economic, environmental or social/cultural impact. Such impacts may include damage to agricultural crops, livestock predation, indigenous fauna predation, soil erosion and land degradation, spread of weeds, pasture/food and habitat competition, and the potential spread of disease. Gong et al. (2009) estimated the overall direct economic impact of several pest animal species (foxes, rabbits, wild dogs, feral pigs, birds and mice) in Australia to be \$740 million annually. This included \$620.8 million of production losses in agriculture (including horticulture) and \$122.7 million on expenditure on pest animal management, administration and research in Australia. Some of the social impacts of pest animals include damage to infrastructure or cultural/historical sites, being 'a nuisance', causing traffic accidents, as well as significant social/psychological impacts on primary producers; for example, through distress of wild dog predation on livestock.

Other non-declared pest animals present in the Mallee include feral Bees, European Wasps, Hares, Starlings, Snails, Rats, Mice and Locusts.

The chenopod dominant vegetation types that dominate the floodplain and those that occupy isolated sandhills on the floodplain have been identified as being susceptible to rabbit activity. Overall, nearly 38,000 hectares (or 80%) of the catchment asset area is considered susceptible to rabbit activity with nearly 8,000 hectares of this being highly susceptible. However, there is little available data to describe the impact rabbits are currently having on the asset area.

The asset area is also known to be home to populations of foxes, pigs, goats and cats. As with rabbits, there is little concrete data available to scale of threat these populations pose this asset.

As with invasive plants, the presence of invasive animals like rabbits and foxes in the asset area is recognised. However, there is only limited data available to adequately describe the current scope and scale of pest animal populations for the purposes of effective regional control. The current information and response capacity relies heavily on local activity at a local scale. This approach, while often very successful, tends to preclude centralised management and control at a regional scale. Therefore, the priority of action to address invasive animals is strongly influenced by local scale data derived from local planning instruments and informal community consultation at the local level. On the other hand, knowledge and understanding of the vulnerability of the Mallee landscape to future incursions or outbreaks is available for some species given the current understanding of their biological requirements and the planning and response framework provided by regional scale plans such as the Mallee Invasive Plants and Animals Management Strategy (IPAWS).

The IPAWS identifies 21 regional assets (based on geographic areas) across the Mallee region. This Catchment Asset area contains four of these geographic assets: Agricultural Land; Tyrrell Creek and Lake Tyrrell; Lalbert Creek and Lake Timboram; and the Southern Buloke Woodland Community. Of these, only the Southern Buloke Woodland Community was considered to be a high priority for onground action to manage invasive plants and animals. The remaining three assets were identified as areas where monitoring for future action and maintaining previous management gains was considered sufficient.

Using this combination of information gathering approaches and the application of a complimentary consultative process, the priority of action for dealing with invasive plants with respect to each Regional Asset has been determined. They are listed in Table 10 below.

Table 10: Prospective Priority of Action to address Invasive Animals upon Regional Assets

Regional Asset	Priority of Action	Summary Logic
Rivers	HIGH	To be documented
Wetlands	HIGH	To be documented
Threatened Species and Communities	HIGH	To be documented

Terrestrial Habitat	HIGH	To be documented
Soils	HIGH	To be documented
Agricultural Land	HIGH	To be documented
Groundwater	NONE	To be documented
Cultural Heritage	HIGH	To be documented
Community Capacity	NONE	To be documented




Figure 13: Susceptibility of Remnant Native Vegetation to Rabbits

Altered Hydrological Regimes

Modification of the natural flow regimes in our river systems has occurred over time to meet the various needs of navigation, agriculture and urban water use. Flow regulation has resulted in changes in the frequency, magnitude and duration of flows, and the restriction of small to medium flood events. River regulation, including the effect of locks, weirs and dams, has altered wetting and drying phases of many wetlands and ephemeral anabranches, by either permanently inundating the area, or restricting flows. Engineering works, such as the building of levees, have alienated large areas of floodplain which alters flood conveyance and flood storage. These changes have great significance for: fringing and floodplain forest communities: populations of fish, macroinvertebrates, algae, macrophytes; nutrient cycling; water quality; and channel shape and form.

Wetlands across the Mallee region have also been subject to modification of natural flow regimes which poses a threat to all priority wetland units. The flow regime, or hydrology, of a wetland is typically determined by climactic conditions and the inflows and outflows of surface and groundwater. Changes in hydrology affect most aspects of wetland ecology, including nutrient cycling; water quality; wetland shape and form; biodiversity; vegetation health, type and extent; and the composition and size of faunal communities. A wetland's flow regime has three main components: frequency, duration and seasonality of inundation. Any activity that changes one or more of these components will alter the natural hydrological regime of a wetland. Examples of these activities include: changes the flow regime of the wetland's water source; interference with flows in and out of, or even within, the wetland; water disposal to or extraction from the wetland; and modification of wetland depth (which alters evaporation rates and affects inundation duration).

The river system adjacent to and anabranch system within this asset area has been subject to extensive modification of hydrological regimes to the influence of weirs and locks and alteration of seasonal flow patterns to benefit irrigated agriculture. Index of Stream Condition analysis in 2004 indicates that, of the 21 river reaches in the asset area, 10 are considered to be in moderate condition and 11 are in a poor condition, primarily due to the typically poor scores for hydrology.

Using this combination of information gathering approaches and the application of a complimentary consultative process, the priority of action for dealing with altered hydrological regimes with respect to each Regional Asset has been determined. They are listed in Table 11 below.

Table 11: Prospective Priority of Action to address Altered Hydrological Processes

Regional Asset	Priority of Action	Summary Logic
Rivers	HIGH	To be documented
Wetlands	HIGH	To be documented
Threatened Species and Communities	HIGH	To be documented
Terrestrial Habitat	HIGH	To be documented
Soils	HIGH	To be documented
Agricultural Land	HIGH	To be documented
Groundwater	NONE	To be documented
Cultural Heritage	HIGH	To be documented
Community Capacity	NONE	To be documented

Soil Erosion

Soil erosion in the Mallee is primarily confined to two processes: wind erosion and water erosion. Wind erosion is typically a regional scale process whereas water erosion primarily occurs in discrete locations on the sides and banks of some watercourses.

Wind erosion is the process by which soil particles are detached from the land surface and transported by the wind. Wind erosion occurs when the forces exerted by the wind overcome the gravitational and cohesive forces of soil particles on the surface of the ground. Wind erosion degrades the soil, reducing its capacity to sustain biodiversity and to support agricultural production. It can also have significant offsite impacts on infrastructure, air quality and respiratory health.

The movement of soil particles by the wind occurs in three ways, each of which is dependent on the size of the particles in question. The first method is known as creep. Larger particles (like sand) are generally too heavy to be lifted by the wind, so they are rolled along the surface by wind drag or moved by bombardment by other moving particles. The next method is known as saltation and this is when middling sized soil particles (typically 0.1 to 0.5 mm) are lifted by the wind, and then fall back to the ground, so they move in a hopping or bouncing fashion. This abrades the soil surface and as they hit other particles they break into smaller particles. It is these smaller particles (typically less than 0.1 mm) that form the suspended dust that is carried away from the erosion site by the wind and is typical of a Mallee dust event.

The susceptibility of a soil to erosion is dependent on its erodibility; its exposure to erosive winds; and on its moisture content. Soil erodibility refers to the inherent properties of the soil that make it susceptible to movement should the soil be exposed to strong winds (for example when they have been completely cleared or cultivated). These properties include: surface texture, organic matter content and stability of

soil aggregates. The most erodible soil types are those with single grained structure and poor aggregate stability, such soils commonly comprising a large proportion of fine sand particles.

Soil erosion susceptibility in the Mallee can be spatially described due to a recognised relationship between soil type and the terrain. Landform component mapping took advantage of the relationship and served as a basis for predicting soil properties that determine erodibility based on prevailing terrain and soil types. It is an extension of the description and mapping of Mallee land systems in 1963 by Rowan and Downes which provides information about the land, its capabilities and its susceptibilities. The mapping used soil texture as the measure of erodibility and considered the frequency and direction of erodible winds (from south-west to westerly) to be consistent across the study area. The erosion susceptibility map for this Catchment Asset area is presented in Figure 14 below.

Wind erosion susceptibility mapping does not reflect the actual threat (or likelihood) of erosion occurring. Whilst a soil's inherent susceptibility to wind erosion is important, the actual likelihood of erosion occurring also depends on how the land is managed. Land management (such as vegetation cover) and climatic conditions (such as wind strength) play an important role in the occurrence of erosion. Where inappropriate management occurs severe wind erosion can be initiated on soils with low susceptibility and conversely erosion can be minimised on soils with high susceptibility with good management practises (Lorimer, 1985). When predicting the threat of erosion, both the landform component's inherent susceptibility to erode and its current state of management must be considered.

Given the heavy clay nature of most soil types within the catchment asset area, very little of the catchment asset is considered to be highly susceptible to wind erosion. Less than 2% of the soils within this asset area are considered to be highly susceptible to wind erosion should the correct precursors (such as reduced groundcover) for such erosion exist. None of the asset area is considered to have very highly susceptible soil types.

Active water erosion of the streamside zone in this asset were identified at 27 discrete locations along the main Murray River channel as part of a river bank erosion survey during 2011/12.

Table 12: Prospective Priority of Action to address Soil Erosion

Regional Asset	Priority of Action	Summary Logic
Rivers	LOW	To be documented
Wetlands	LOW	To be documented
Threatened Species and Communities	LOW	To be documented
Terrestrial Habitat	LOW	To be documented
Soils	LOW	To be documented
Agricultural Land	LOW	To be documented
Groundwater	NONE	To be documented
Cultural Heritage	HIGH	To be documented
Community Capacity	NONE	To be documented




Figure 14: Wind Erosion Susceptibility

Inappropriate Water Use Practices

Inappropriate water use practices in agricultural activities in both the irrigation and dryland zones have been demonstrated to result in excessive volumes of deep drainage past the root zone of crops and pasture and therefore contribute to the raising of local and regional water tables.

Approximately 93% (over 350,000 hectares) of the asset area is given over to dryland agricultural activities. There is insufficient information to determine the extent to which inappropriate water use practises remain in place.

This catchment asset contains only a relatively small area of agricultural activities compared to many other asset areas. Approximately 10% of the asset area is given over to agricultural activities – almost 1,700 hectares of irrigation and a further 3,200 hectares of dryland farming. There is a large area of dryland agriculture to the east of Lake Cullulleraine that contains a mix of cropping and grazing activities. Lindsay Point in the far west of the asset area is the only significant irrigation area within the asset. There is insufficient information to determine the extent to which inappropriate water use practises remain in place in either agricultural area.

Table 13: Prospective Priority of Action to address Inappropriate Water Use Practices

Regional Asset	Priority of Action	Summary Logic
Rivers	HIGH	To be documented
Wetlands	HIGH	To be documented
Threatened Species and Communities	HIGH	To be documented
Terrestrial Habitat	HIGH	To be documented
Soils	HIGH	To be documented
Agricultural Land	HIGH	To be documented
Groundwater	NONE	To be documented
Cultural Heritage	HIGH	To be documented
Community Capacity	NONE	To be documented

Recreational Pressures

Recreational pressure can contribute to impacts including littering, track proliferation, fishing pressures, firewood collection, soil compaction and site erosion. The nature of the impacts is typically localised around the particular site and is highly dependent on the accessibility, popularity and sensitivity of the site along with the level of management that the location receives.

This catchment asset contains some of the most popular areas for recreation activities within this region. Lindsay Island, Wallpolla Island and Merbein Common are some of the most popular and accessible riverine areas though there are many others. The State of the Parks survey identifies a range of potential impacts arising from this recreational use. However, the scope, scale and extent of the impacts that results from this recreational pressure have not been quantified.

Table 14: Prospective Priority of Action to address Recreational Pressures

Regional Asset	Priority of Action	Summary Logic
Rivers	HIGH	To be documented
Wetlands	HIGH	To be documented
Threatened Species and Communities	HIGH	To be documented
Terrestrial Habitat	HIGH	To be documented
Soils	NONE	To be documented
Agricultural Land	NONE	To be documented
Groundwater	NONE	To be documented
Cultural Heritage	HIGH	To be documented
Community Capacity	NONE	To be documented

Land Use Change

Land use change as a threatening process in this context is considered to be the change of land management or use practices from either a steady state or from accepted best practice management system. Examples include the removal of native vegetation, conversion of dryland property to irrigation development (or the reverse), change from no-till cropping to traditional fallow techniques, conversion from perennial pasture to cropping (or the reverse)

There is insufficient quality and quantity of data available to adequately determine the nature and level of impact resulting from this threatening process on this catchment asset at this time.

Table 15: Prospective Priority of Action to address Land Use Change

Regional Asset	Priority of Action	Summary Logic
Rivers	HIGH	To be documented
Wetlands	HIGH	To be documented
Threatened Species and Communities	HIGH	To be documented
Terrestrial Habitat	HIGH	To be documented
Soils	HIGH	To be documented
Agricultural Land	HIGH	To be documented

Groundwater	NONE	To be documented
Cultural Heritage	HIGH	To be documented
Community Capacity	NONE	To be documented

Direct Off-site Interactions

Direct physical impacts from land management activities on neighbouring off-site assets such as areas of remnant native vegetation or wetlands. Such interactions may include chemical spray drift; parking or storage of machinery and equipment; or incremental drift of cultivation into the asset.

There is insufficient quality and quantity of data available to adequately determine the nature and level of impact resulting from this threatening process on this catchment asset at this time.

Table 16: Prospective Priority of Action to address Direct Off-site Interactions

Regional Asset	Priority of Action	Summary Logic
Rivers	LOW	To be documented
Wetlands	LOW	To be documented
Threatened Species and Communities	LOW	To be documented
Terrestrial Habitat	LOW	To be documented
Soils	LOW	To be documented
Agricultural Land	LOW	To be documented
Groundwater	NONE	To be documented
Cultural Heritage	LOW	To be documented
Community Capacity	NONE	To be documented

Misaligned Community Perceptions

Community opinions, approaches & values that run counter to the messages & knowledge available about natural resource management AND threaten the success of the wider communities' efforts to enhance their environment. Such perceptions include 'right of unfettered access' that results in removal of traffic management infrastructure installed near river banks; and 'we are doing no harm' where individuals are not aware of the cumulative and incremental harm of some of their actions (along with those of the rest of the community) when they are making use of our assets

There is insufficient quality and quantity of data available to adequately determine the nature and level of impact resulting from this threatening process on this catchment asset at this time. However, there is a range of anecdotal information from a range of sources that suggest the threat is present and ongoing.

Table 17: Prospective Priority of Action to address Misaligned Community Perceptions

Regional Asset	Priority of Action	Summary Logic
Rivers	MEDIUM	To be documented
Wetlands	MEDIUM	To be documented
Threatened Species and Communities	MEDIUM	To be documented
Terrestrial Habitat	MEDIUM	To be documented

Soils	MEDIUM	To be documented
Agricultural Land	MEDIUM	To be documented
Groundwater	NONE	To be documented
Cultural Heritage	MEDIUM	To be documented
Community Capacity	MEDIUM	To be documented

Inappropriate Fire Regimes

Fire is an ongoing challenge for land managers and communities alike. Fire is also a major force determining the structure, function and sustainability of Australia's ecosystems. A substantial proportion of Australia's unique biota is dependent, to varying degrees, on fire and the variety of fire regimes for its continued existence and development. In this context, inappropriate fire regimes can mean either too little or too much fire

There is insufficient quality and quantity of data available to adequately determine the nature and level of impact resulting from this threatening process on this catchment asset at this time.

Table 18: Prospective Priority of Action to address Inappropriate Fire Regimes

Regional Asset	Priority of Action	Summary Logic
Rivers	NONE	To be documented
Wetlands	NONE	To be documented
Threatened Species and Communities	LOW	To be documented
Terrestrial Habitat	LOW	To be documented
Soils	NONE	To be documented
Agricultural Land	NONE	To be documented
Groundwater	NONE	To be documented
Cultural Heritage	NONE	To be documented
Community Capacity	NONE	To be documented




Figure 15: Fire History

Constrained Regenerative Capacity

The decline in vegetation cover and habitat complexity within remnant native vegetation can constrain or prevent regeneration which can lead to loss of habitat in the longer term. There are many contributors to this threatening process including weed invasion, excess grazing pressure, and habitat fragmentation. Loss of understorey flora and associated fauna are a possible outcome, also leading to a reduction in the capacity of the remnant to support flora and fauna species or maintain current population numbers, thus impacting on the biodiversity value of the asset.

Habitat fragmentation is one useful indicator of constrained regenerative capacity. A high level of fragmentation will severely limit the capacity of a vegetation community and the ecology it supports to maintain its health and reproductive capacity. It will also hamper our efforts to improve the health and condition of remnant landscapes at a broad scale due to the level of additional resources that each remnant patch would require (such as restoration or revegetation) rather than being able to simply protect the remnant (through fencing and invasive species control) and then rely upon its own internal regenerative resources to bring about an improvement in health and condition.

Modelling of habitat fragmentation using a spatial tool (Vogt, et al. 2007) can demonstrate the level of habitat fragmentation of a landscape. Using a combination of information gathering approaches and the application of a complimentary consultative process, the priority of action for dealing with constrained regenerative capacity with respect to each Regional Asset has been determined. They are listed in Table 19 below.

Table 19: Prospective Priority of Action to address Constrained Regenerative Capacity

Regional Asset	Priority of Action	Summary Logic
Rivers	MEDIUM	To be documented
Wetlands	MEDIUM	To be documented
Threatened Species and Communities	MEDIUM	To be documented
Terrestrial Habitat	MEDIUM	To be documented
Soils	NONE	To be documented

Agricultural Land	NONE	To be documented
Groundwater	NONE	To be documented
Cultural Heritage	NONE	To be documented
Community Capacity	NONE	To be documented




Figure 16: Habitat Fragmentation

Section 4 – References & Appendices

References

Department of Agriculture, Fisheries and Forestry. *National Natural Resource Management Capacity Building Framework*. Canberra: Department of Agriculture, Fisheries and Forestry, 2005.

Gong, W., J. Sinden, M. Braysher, and R. Jones. *The economic impacts of vertebrate pests in Australia*. Canberra: Invasive Animals Cooperative Research Centre, 2009, 59.

Lorimer, M. S. "Estimating the susceptibility of soil to wind erosion in Victoria." *Journal of Australian Institute for Agricultural Science*, *51* (2), 1985: 122-127.

Mock, I., and V. Grinter. *Mapping the Mallee's Saline Land. Stage 3: Classifying Mapped Salinity. Final Report*, Irymple: Mallee Catchment Management Authority, 2009.

Vogt, P., K. Riitters, C. Estreguil, J. Kozak, T. Wade, and J. Wickham. "Mapping spatial patterns with morphological image processing." *Landscape Ecology* 22, 2007: 171-177.

The Value of Regional Assets within a Catchment Asset

The Regional Asset value indicates the relative importance of the Regional Asset within the Catchment Asset Area. The valuation is determined from a range of indicators, many specific to each of the Regional Assets, that describe the importance of the Asset relative to social, economic and environmental values. One common example of an indicator of asset value is legislative significance which provides an indication of whether part or all of a Regional Asset is of significance at a local, regional, State, National or international level. These indicators were grouped and scored by consensus against agreed criteria and then each Regional Asset within the Catchment Asset was assigned a category that defines the prospective asset value and relative importance of and priority for maintaining or improving the value of the Asset.

Five category levels were identified:

None – indicates that the Regional Asset is unknown or not present within the Catchment Asset area.

Low – indicates that the Regional Asset is of relatively low status or importance within the Catchment Asset area. It may also indicate that there is insufficient knowledge or data available to define or quantify the asset value and therefore prioritise it with confidence. Further research may be required to rectify this situation.

Medium – indicates that the Regional Asset is locally or regionally important but relatively well known or secure at the Catchment Asset scale. This may be due in part to previously successful management interventions. Ongoing monitoring may be required as will some level of intervention (particularly at a local scale) to ensure previous management gains are not lost.

High – indicates that the Regional Asset within the Catchment Asset Area is of significance at a regional, State or National scale and that action is likely required over the life of the RCS in order to maintain or improve the value of the Regional Asset within the Catchment Asset Area.

Very High - indicates that the Regional Asset within the Catchment Asset Area is of significance at a National or international scale and that action is likely required over the life of the RCS in order to maintain or improve the value of the Regional Asset within the Catchment Asset Area.

2. Categorising Threatening Processes

The categorisation of the threatening processes in order to prioritise management actions was based on the current scope and scale of the impacts arising from the threatening process along with consideration of potential future impact of the threatening process. The categorisation was based on consensus against agreed criteria. Each threatening process acting upon each Regional Asset was assigned a category that defines the prospective priority of action to address that threatening process.

Four category levels were identified:

None – indicates that either the threatening process, or the regional asset itself, is unknown or not present within the catchment Asset area.

Low – indicates that, while the threatening process is known to be present, it is at a low level or the Regional Asset itself is not overly susceptible to or impacted by the threat. It may also indicate that there is insufficient knowledge or data available to define or quantify the interaction between the asset and the threatening process and therefore prioritise it with confidence. Further research may be required to rectify this situation.

Medium – indicates that the threatening process is real but that the Regional Asset is not under immediate threat at the Catchment Asset scale. This may be due to previously successful interventions.

Ongoing monitoring will be required as will some level of intervention (particularly at a local scale) to ensure previous management gains are not lost.

High – indicates that: there is a high level of potential harm to the Asset as a result of the threatening process; there are a variety of management actions available that are known to be successful at reducing harm and which are generally supported by land managers; and that action is required over the life of the RCS in order to prevent further impact as a result of the threatening process.

3. Management Plans Relevant to this Catchment Asset

Table 20: Management Plans Relevant to this Catchment Asset

Name	Туре	Author/Owner	Year
North West Region Mildura Fire District Fire Protection Plan	Management Plan	Department of Sustainability & Environment	2008
Merbein to SA Border Murray River Frontage Action Plan	Management Plan	Mallee Catchment Management Authority	2003
Victorian Mallee Irrigation Area Land and Water Management Plan (Draft)	Management Plan	Mallee Catchment Management Authority	2011
The River Murray Channel Icon Site Environmental Management Plan 2006-2007	Management Plan	Murray Darling Basin Authority	2006
Lindsay-Walpolla Islands Environmental Water Management Plan 2012	Management Plan	Murray Darling Basin Authority	2012
Mallee Parks Management Plan	Management Plan	Parks Victoria	1996
Northern Region Sustainable Water Strategy	Strategy	Department of Sustainability & Environment	2009
Mallee River Health Strategy 2005	Strategy	Mallee Catchment Management Authority	2006
Mallee Wetland Strategy 2006-2011	Strategy	Mallee Catchment Management Authority	2006
Native Fish Strategy for the Murray-Darling Basin 2003-2013	Strategy	Murray Darling Basin Authority	2003
Mallee Invasive Plants and Animals Management Strategy	Strategy	Mallee Catchment Management Authority	2011
Millewa Carwarp Landcare Group Action Plan 2012-2017	Action Plan	Millewa Carwarp Landcare Group	2012
National Recovery Plan for the Murray Cod Maccullochella peelii peelii	National Recovery Plan	Department of Sustainability & Environment (Victoria)	2010
National Recovery Plan for the Murray Hardyhead Craterocephalus fluviatilis	National Recovery Plan	Department of Sustainability & Environment (Victoria)	2008
National Recovery Plan for the Regent Parrot (eastern subspecies) Polytelis anthopeplus monarchoides	National Recovery Plan	Department of Sustainability & Environment (Victoria)	2011

		T	
Flora & Fauna Guarantee Action Statement Plaucident (Giles) Planigale Planigale gilesi	Action Statement	Department of Sustainability & Environment (Victoria)	2003
Flora & Fauna Guarantee Action Statement White-bellied Sea-eagle Haliaeetus leucogaster	Action Statement	Department of Sustainability & Environment (Victoria)	2003
Flora & Fauna Guarantee Action Statement Spotted Bowerbird Chlamydera maculata	Action Statement	Department of Sustainability & Environment (Victoria)	2003
Flora & Fauna Guarantee Action Statement White-browed Treecreeper Climacteris affinis	Action Statement	Department of Sustainability & Environment (Victoria)	2003
Flora & Fauna Guarantee Action Statement Bush Stone-Curlew Burhinus grallarius	Action Statement	Department of Sustainability & Environment (Victoria)	2004
Flora & Fauna Guarantee Action Statement Grey Falcon Falco hypoleucos	Action Statement	Department of Sustainability & Environment (Victoria)	2003
Flora & Fauna Guarantee Action Statement Major Mitchell's Cockatoo Cacatua leadbeateri	Action Statement	Department of Sustainability & Environment (Victoria)	2004
Flora & Fauna Guarantee Action Statement Freckled Duck Stictonetta naevosa	Action Statement	Department of Sustainability & Environment (Victoria)	2004
Flora & Fauna Guarantee Action Statement Hooded Scaly-foot Pygopus nigriceps	Action Statement	Department of Sustainability & Environment (Victoria)	2004
Flora & Fauna Guarantee Action Statement Barking Owl Ninox connivens	Action Statement	Department of Sustainability & Environment (Victoria)	2004
Flora & Fauna Guarantee Action Statement Great Egret Ardea alba Intermediate Egret Ardea intermedia Little Egret Egretta garzetta	Action Statement	Department of Sustainability & Environment (Victoria)	2001
Flora & Fauna Guarantee Action Statement Twelve threatened Swainson-peas and Darling-peas (Swainsona species)	Action Statement	Department of Sustainability & Environment (Victoria)	2003
Flora & Fauna Guarantee Action Statement Blue-billed Duck Oxyura australis	Action Statement	Department of Sustainability & Environment (Victoria)	2003
Flora & Fauna Guarantee Action Statement Inland Carpet Python Morelia spilota metcalfei	Action Statement	Department of Sustainability & Environment (Victoria)	2003
Flora & Fauna Guarantee Action Statement Painted Honeyeater Grantiella picta	Action Statement	Department of Sustainability & Environment (Victoria)	2003
Flora & Fauna Guarantee Action Statement Freshwater Catfish Tandanus tandanus	Action Statement	Department of Sustainability & Environment (Victoria)	2005

Flora & Fauna Guarantee Action Statement Silver Perch Bidyanus bidyanus	Action Statement	Department of Sustainability & Environment (Victoria)	2005
---	------------------	---	------

4. Management Action Definitions

Management Action Category	Management Action	Definition	Threatening Processes Addressed
On-ground Works	Pest Plant Control	Control and eradication of pest plant species	Invasive Plants; Constrained Regenerative Capacity
	Pest Animal Control	Control and eradication of pest animal species	Invasive Animals; Soil Erosion; Constrained Regenerative Capacity
	Habitat Protection	Habitat protection through exclusion fencing or other physical means	Invasive Animals; Soil Erosion; Recreational Pressure; Land Use Change; Direct Off-Site Interactions; Constrained Regenerative Capacity
	Habitat Restoration	Restoration of degraded terrestrial habitat by planting vegetation	Land & Water Salinisation; Soil Erosion; Recreational Pressure; Direct Off-Site Interactions; Constrained Regenerative Capacity
	Revegetation	Re-creation of terrestrial habitat by planting vegetation (e.g. new wildlife corridor plantings)	Land & Water Salinisation; Soil Erosion; Recreational Pressure; Direct Off-Site Interactions; Constrained Regenerative Capacity
	Environmental Watering	Delivering environmental water to wetlands and floodplains; maintaining appropriate watering regimes	Altered Hydrological Regimes; Constrained Regenerative Capacity
	Soil Erosion Control	Control of soil erosion through engineering works and other structures	Soil Erosion
	Threatened Species Interventions	Interventions to improve outcomes for threatened species and ecological communities	Constrained Regenerative Capacity
	Enhancing Land Management Regimes	Interventions to enhance land management regimes on both public and private land (e.g. water use efficiency on irrigated land; maintaining appropriate soil cover in dryland agriculture; improved management or protection regimes)	Land & Water Salinisation; Soil Erosion; Inappropriate Water Use Practises; Land Use Change; Direct Off-Site Interactions
Capacity Building	Supporting Human Capacity for NRM	Provision of new skills and training in NRM; Delivery of publicity and awareness raising paraphernalia	All
	Supporting Institutional Capacity for NRM	Development and support of regional partnerships between institutions with a stake in NRM	All
	Supporting Social Capacity for NRM	Supporting community organisations that have a stake in NRM by aiding	All

		governance and provision of material	
NRM Planning	Institutional Planning for NRM	Development and implementation of NRM planning by regional institutions. Examples include institutional NRM plans, regional asset plans and substrategies	All
	Community Driven Planning for NRM	Development and implementation of NRM planning by community based organisations. Examples include Landcare Group Action Plans.	All
	Landholder Driven Planning for NRM	Development and implementation of NRM planning by landholders. Examples include property management plans and conservation agreements	All
Knowledge Building	Research to improve knowledge	Any research or other work delivered to improve or expand current knowledge or fill knowledge gaps in any NRM topic relevant to regional assets	All
Asset Assessment	Asset condition monitoring and assessment	Monitoring and collection of data concerning indicators of regional asset condition and/or threatening processes impacting on regional assets; Efforts to determine and/or evaluate trends in the condition of assets or the scope and scale of threatening processes impacting on regional assets.	All

5. Management Actions from Existing Management Plans

Table 21: Management Actions from Existing Management Plans

Instrument	Management Action	Details
North West Region Mildura Fire District	Supporting Institutional Capacity for NRM	Investing in integrated fire management to balance community needs with ecological requirements
Fire Protection Plan	Research to improve knowledge	Determining appropriate ecological fire regimes
Merbein to SA Border	Pest Plant Control	reducing the impact of pest plants on frontages
Murray River Frontage Action Plan	Pest Animal Control	reducing the impact of pest animals on frontages
	Habitat Protection	definition of camp sites using vehicle barriers; track rationalisation; stock containment fencing; protect frontages and instream habitat with high biodiversity values; protect cultural heritage sites

	Habitat Restoration	Restore degraded frontages
	Revegetation	Restore degraded frontages
	Environmental Watering	Use watering to restore degraded frontages
	Supporting Human Capacity for NRM	education campaign for recreational users, grazing license holders; cultural heritage education activities
	Supporting Institutional Capacity for NRM	capacity raising and partnerships between regional authorities; Coordinate early detection and rapid control responses for new and emerging weeds
	Research to improve knowledge	R&D to improve knowledge of floodplain nutrient dynamics and effectiveness of measures to minimise sedimentation and nutrient inputs to Murray River
	Asset condition monitoring and assessment	Monitor total grazing pressures
Victorian Mallee Irrigation Area Land and Water	Enhancing Land Management Regimes	improving water use efficiency; modernising infrastructure
Management Plan (Draft)	Supporting Human Capacity for NRM	Education and community awareness development on a range of land, water and biodiversity management issues
	Supporting Institutional Capacity for NRM	Maintaining partnerships and communication channels amongst regional and State partners
	Institutional Planning for NRM	Considerable (and diverse) planning and policy development to enhance water trading rules, improve efficiency of water use and incorporate environmental watering into the water management process
	Landholder Driven Planning for NRM	Whole farm planning for irrigation properties
	Research to improve knowledge	Various research topics on water use efficiency improvements and the potential impacts of irrigation on biodiversity and land use management
	Asset condition monitoring and assessment	waterway salinity, groundwater and irrigation drainage monitoring
The River Murray	Pest Plant Control	Invasive species management
Channel Icon Site Environmental	Pest Animal Control	Invasive species management
Management Plan 2006-2007	Habitat Restoration	restoration of fish passage through locks and weirs
	Environmental Watering	Regulating structures for specific wetlands and anabranches (exclude un-seasonal flows; improve wetting/drying regime)
Lindsay-Wallpolla Islands Environmental Water Management Plan 2012	Environmental Watering	Define and/or apply ecological watering requirements of antecedent ecological systems
Mallee Parks Management Plan	Pest Plant Control	reducing the impact of pest plants on native species and communities
	Pest Animal Control	reducing the impact of pest animals on native species and communities

Habitat Protection the existing diversity of native flora and fauna w maintained, and a management regime of minir disturbance adopted. Significant natural and cu features will be given special protection	
reatares will be given epostal protestion	
Habitat Restoration active revegetation in areas of localised extincti rarity; adopting manipulative fire regimes where can be demonstrated to be of value to the Park environments and ecosystems	they
Revegetation Wildlife corridors between the Parks and other blocks of public land in the region will be established.	
Environmental Watering Restoring a more natural hydrological regime w Parks	ithin all
Threatened Species Manage Flora and Fauna Guarantee listed species communities according to approved action states	
Supporting Human Capacity for NRM Mumerous actions to assist visitors to discover, and appreciate the natural and cultural features Mallee Parks, and the value of National Parks a management	of the
Supporting Institutional Capacity for NRM Liaise with local community groups and land ho and, as appropriate, involve them in relevant as of planning and managing the Parks	
Research to improve knowledge Encourage research on Mallee ecosystems, ve communities, and significant flora and fauna sp	
Asset condition Monitor control and rehabilitation programs; Su and monitor vegetation and habitats on Lindsay assessment	
Northern Region Sustainable Water Environmental Watering A range of policies and actions at a regional scar are intended to support environmental water ac	
Strategy Supporting Institutional Capacity for NRM A range of policies and actions that support inst partnership arrangements to improve bulk wate management for irrigators, consumers and the environment	
Asset condition Water quality and quantity monitoring in surface groundwater resources; ecological monitoring of assessment and wetlands	
Mallee River Health Strategy 2005Pest Plant Control weed control programsPlanning, coordination and implementation of p weed control programs	riority
Pest Animal Control Planning, coordination and implementation of p pest control programs	riority
Supporting Institutional Capacity for NRM Partnerships between State and regional organ to close knowledge gaps and provide for co-ord implementation of a range of management plan relevant to this catchment asset	linated
Supporting Social Community education via schools, Landcare an Waterwatch groups	ıd
Research to improve knowledge Numerous research topics to improve knowledge river reaches and their ecological requirements this catchment asset	
Pest Plant Control Focus on willows and aquatic weeds	

Mallee Wetland	Pest Animal Control	Focus on rabbits & carp
Strategy 2006-2011	Habitat Protection	Identification of priority sites; protection from stock grazing
	Habitat Restoration	Identification of priority sites with works to occur at five of them across the region
	Environmental Watering	Determine appropriate water regimes
	Threatened Species Interventions	Identify three priority species and implement key recommendations from their Action Statements or Recovery Plans as relevant to wetlands
	Supporting Human Capacity for NRM	Raising awareness and improving knowledge of wetland values and threats in the community; improving knowledge of indigenous wetland values
	Institutional Planning for NRM	preparation and implementation of local planning overlays and wetland management plans
	Research to improve knowledge	improve knowledge of wetland values and threats to those values
	Asset condition monitoring and assessment	Determine wetland condition; monitor responses arising from interventions
Native Fish Strategy for the Murray-Darling	Pest Animal Control	Control of carp and other alien fish
Basin 2003-2013	Habitat Protection	protect key aquatic and riparian habitat; protect key linkages between floodplains and aquatic habitats
	Habitat Restoration	improve fish passage/water quality/key aquatic & riparian habitats
	Institutional Planning for NRM	preparation and implementation of species and fish community recovery plans
	Supporting Institutional Capacity for NRM	enhance knowledge of and partnership support for the Strategy
	Supporting Human Capacity for NRM	enhance knowledge of and support for the Strategy and native fish in general
Mallee Invasive Plants and Animals	Pest Plant Control	prevent new introductions; contain high risk weed species
Management Strategy 2011	Pest Animal Control	prevent new introductions; contain high risk pest animal species
	Supporting Human Capacity for NRM	improve community knowledge and awareness of invasive plants and animals and methods of control
	Supporting Institutional Capacity for NRM	support partnership arrangements between organisations with responsibilities for control and management of invasive plants and animals
	Supporting Social Capacity for NRM	Support community organisations carrying out targeted invasive plant and animal management with information, education, extension, enforcement and the identification of funding opportunities; Actively seek community participation and create partnerships with community groups in all IPA management project and programs
	Research to improve knowledge	Investigate status and impacts arising from fox predation on turtles

	Asset condition monitoring and	monitor invasive plants and animals against both past interventions and for future actions;
	assessment	,
National Recovery Plan for the Murray Cod Maccullochella	Environmental Watering	Develop and implement flow management practices to benefit recovery of Murray Cod populations
peelii peelii	Supporting Human Capacity for NRM	
	Supporting Institutional Capacity for NRM	
	Research to improve knowledge	
	Asset condition monitoring and assessment	
National Recovery Plan for the Murray Hardyhead Craterocephalus	Threatened Species Interventions	Establish and maintain the Murray Hardyhead in captivity; Establish new populations of Murray Hardyhead in the wild
fluviatilis	Supporting Human Capacity for NRM	
	Research to improve knowledge	
	Asset condition monitoring and assessment	
National Recovery	Revegetation	filling gaps in and expanding width of known flyways
Plan for the Regent Parrot (eastern subspecies) Polytelis	Threatened Species Interventions	
anthopeplus monarchoides	Supporting Human Capacity for NRM	Increase community involvement in the eastern Regent Parrot recovery program
	Supporting Institutional Capacity for NRM	
	Institutional Planning for NRM	
	Research to improve knowledge	
	Asset condition monitoring and assessment	
Flora & Fauna Guarantee Action	Institutional Planning for NRM	update land management plans to reflect habitat requirements for species
Statement Plaucident (Giles) Planigale Planigale gilesi	Research to improve knowledge	
Flora & Fauna	Habitat Protection	buffers/barriers to protect nest sites from disturbance
Guarantee Action Statement White- bellied Sea-eagle	Supporting Human Capacity for NRM	community education
Haliaeetus leucogaster	Institutional Planning for NRM	incorporate suitable habitat into appropriate management plans in order to protect such habitat

	l					
	Research to improve knowledge	improve knowledge of ecological requirements/potential habitat/recovery techniques				
	Asset condition monitoring and assessment	Species population monitoring; threat status monitoring				
Flora & Fauna Guarantee Action	Supporting Human Capacity for NRM	Extension and awareness raising to raise profile of species				
Statement Spotted Bowerbird Chlamydera maculata	Asset condition monitoring and assessment	Monitor for presence in catchment asset				
Flora & Fauna	Pest Animal Control	rabbit control in pine-buloke woodlands				
Guarantee Action Statement White- browed Treecreeper	Habitat Restoration	restoration of degraded pine-buloke woodland core species habitat				
Climacteris affinis	Supporting Human Capacity for NRM	training and extension to improve staff and community identification of species				
	Research to improve knowledge	improve knowledge of ecological requirements/potential habitat/recovery techniques				
Flora & Fauna Guarantee Action	Pest Animal Control	Fox control in and near key habitat				
Statement Bush Stone-Curlew Burhinus grallarius	Supporting Human Capacity for NRM	Raise community awareness of species; target golf courses with remnant habitat with specific education and land management material				
	Institutional Planning for NRM	Develop or amend planning scheme regulations to protect key habitat areas				
	Landholder Driven Planning for NRM	Encourage and support a) fencing of remnant habitat; and/or b) the creation of covenants; to protect species on private land				
Flora & Fauna Guarantee Action	Supporting Human Capacity for NRM	community education				
Statement Grey Falcon Falco hypoleucos	Research to improve knowledge	Numerous research topics to improve knowledge on species and its requirements				
Flora & Fauna Guarantee Action	Habitat Restoration	conserve and enhance species preferred habitat				
Statement Major Mitchell's Cockatoo Cacatua leadbeateri	Supporting Human Capacity for NRM	Raise community awareness of species; training and extension to improve staff and community knowledge of species requirements				
	Research to improve knowledge	improve knowledge of ecological requirements/potential habitat/recovery techniques				
Flora & Fauna Guarantee Action	Threatened Species Interventions	wetland closures to hunting				
Statement Freckled Duck Stictonetta naevosa	Supporting Human Capacity for NRM	hunter education				
	Research to improve knowledge	improve knowledge of ecological requirements/potential habitat/recovery techniques				
	Asset condition monitoring and assessment	targeted searches				
Flora & Fauna Guarantee Action	Research to improve knowledge	improve knowledge of ecological requirements/potential habitat/recovery techniques				

Statement Hooded Scaly-foot Pygopus nigriceps	Asset condition monitoring and assessment	targeted searches					
Flora & Fauna Guarantee Action	Supporting Human Capacity for NRM	Raise community awareness of species					
Statement Barking Owl Ninox connivens	Supporting Institutional Capacity for NRM	Support municipal councils and other agencies responsible for land management in effectively managing species habitat					
	Institutional Planning for NRM	Alter and update appropriate plans to reflect species ecology and habitat requirements					
	Landholder Driven Planning for NRM	Encourage and support the creation of covenants to protect species on private land					
	Research to improve knowledge	improve knowledge of ecological requirements/potential habitat/recovery techniques					
	Asset condition monitoring and assessment	targeted searches					
Flora & Fauna Guarantee Action	Environmental Watering	Maintain appropriate water regimes in wetlands and creeks systems					
Statement Great Egret Ardea alba Intermediate Egret Ardea intermedia Little Egret Egretta garzetta	Asset condition monitoring and assessment	Species population monitoring; threat status monitoring					
Flora & Fauna Guarantee Action	Pest Plant Control	Control competing weeds in known populations					
Statement Twelve threatened Swainson-	Habitat Protection	Exclusion of stock and vehicular access to known habitat/populations Raise community awareness of species					
peas and Darling- peas (Swainsona species)	Supporting Human Capacity for NRM						
	Institutional Planning for NRM	prepare and implement management plans for public reserves containing species					
	Landholder Driven Planning for NRM	Encourage and support a) fencing of remnant habitat; and/or b) the creation of covenants; to protect species on private land					
	Research to improve knowledge	improve knowledge of ecological requirements/potential habitat/recovery techniques					
	Asset condition monitoring and assessment	Species population monitoring; threat status monitoring					
Flora & Fauna	Pest Animal Control	Fox control in and near key habitat					
Guarantee Action Statement Blue-billed Duck Oxyura australis	Habitat Protection	Exclusion of stock and vehicular access to known habitat/populations; minimise disturbance through wetland closure etc.					
	Supporting Human Capacity for NRM	Raise community awareness of species; target golf courses with remnant habitat with specific education and land management material					

	I					
	Landholder Driven Planning for NRM	Encourage and support a) fencing of remnant habitat; and/or b) the creation of covenants; to protect species on private land				
	Research to improve knowledge	improve knowledge of ecological requirements/potential habitat/recovery techniques				
	Asset condition monitoring and assessment	Species population monitoring; threat status monitoring				
Flora & Fauna Guarantee Action	Habitat Protection	Exclusion of stock and vehicular access to habitat; management of recreation activities in preferred habitat				
Statement Inland Carpet Python	Pest Animal Control	Rabbit and fox control				
Morelia spilota metcalfei	Threatened Species Interventions	Numerous interventions to manage known populations				
	Supporting Human Capacity for NRM	Raise community awareness of species				
	Research to improve knowledge	Numerous research topics to improve knowledge on species and its requirements				
	Asset condition monitoring and assessment	Species population monitoring				
Flora & Fauna Guarantee Action	Supporting Human Capacity for NRM	awareness raising to improve reporting rates				
Statement Painted Honeyeater Grantiella picta	Institutional Planning for NRM	update regional planning instrument to reflect species habitat requirements				
	Research to improve knowledge	encourage university and other institutions to conduct appropriate ecological research				
	Asset condition monitoring and assessment	survey known habitats				
Flora & Fauna Guarantee Action	Environmental Watering	provide for appropriate water regimes to suit species requirements				
Statement Freshwater Catfish Tandanus tandanus	Threatened Species Interventions	reintroductions				
	Supporting Human Capacity for NRM	awareness raising of species status and means to minimise harm to species				
	Institutional Planning for NRM	planning and prioritisation tools developed or modified				
	Research to improve knowledge	improve knowledge of ecological requirements/potential habitat/recovery techniques				
	Asset condition monitoring and assessment	routine species population survey				
Flora & Fauna Guarantee Action	Environmental Watering	provide for appropriate water regimes to suit species requirements				
Statement Silver	Threatened Species Interventions	reintroductions				

Perch Bidyanus bidyanus	Supporting Human Capacity for NRM	awareness raising of species status and means to minimise harm to species			
	Institutional Planning for NRM	planning and prioritisation tools developed or modified			
	Research to improve knowledge	improve knowledge of ecological requirements/potential habitat/recovery techniques			
	Asset condition monitoring and assessment	routine species population survey			
Millewa Carwarp	Pest Animal Control	rabbit control			
Landcare Group Action Plan 2012-2017	Supporting Institutional Capacity for NRM	improve group governance; develop group capacity; partnership with TfN			
	Supporting Social Capacity for NRM	support for Junior Landcare			
	Community Driven Planning for NRM	identify and plan for biodiversity assets within group area			

6. Prioritising Potential Management Actions

Potential management action categories are prioritised in terms of:

- the value of a regional asset within the catchment asset area;
- the priority to address a threatening process to those values;
- the capacity of the management action to address the threatening process; and
- the level of investment (in both of time and money) required to exercise that capacity.

Individual management actions are grouped and scored by consensus against agreed criteria and then each potential management action group within the Catchment Asset is assigned to a category that defines the prospective priority to the implementation.

Four category levels were identified:

None – indicates that either the management action group not contained in management plans relevant to the Catchment Asset area or the targeted threatening processes that it addresses is unknown or not present within the Catchment Asset area.

Low – typically indicates either that the targeted threatening processes are of low priority within this Catchment Asset area or the management action category is known to be generally not effective at mitigating the threatening process. It may also indicate that there is insufficient knowledge or data available to define or quantify the effectiveness of the management action to mitigate the threatening process and therefore prioritise it with confidence. Further research may be required to rectify this situation.

Medium – typically indicates that the targeted threatening processes are of medium priority within this Catchment Asset area and the management action category is known to be generally effective at mitigating the threatening process and is within the region's capacity to implement over the life of the RCS. It may also indicate that there is some uncertainty with the effectiveness of the potential management action upon a high priority threatening process in terms of either the scale of the threat, the known capacity of the action or the investment required to exercise that capacity.

High – indicates that the targeted threatening processes are of high priority within this Catchment Asset area and the management action category is known to be effective at mitigating the threatening process and is within the region's capacity to implement over the life of the RCS and that action is required urgently in order to prevent further impact as a result of the threatening process.

7. Management Action Priorities

Threatening Processes Priority of Action

Table 22 below represents the findings of a group based analysis in June 2103 of the threat level posed by threatening processes upon Regional Assets within this Catchment Asset based on the criteria contained in Appendix 2. *Categorising Threatening Processes*. The bottom row of the table contains a median 'score' for the threat level posed by each of the threatening processes across the Catchment Asset. These 'scores' are the same as those shown in Table 2.

Table 22: Threatening Process Priority of Action Summary

	Land & water salinisation	Invasive Plants	Invasive Animals	Altered Hydrological Regimes	Soil Erosion	Inappropriate Water Use Practices	Recreational Pressures	Land Use Change	Direct off-site interactions	Misaligned community perceptions	Inappropriate fire regimes	Constrained regenerative capacity
Rivers	High	High	High	High	Medium	High	High	High	Low	Medium	None	High
Wetlands	High	High	High	High	Medium	High	High	High	Low	Medium	None	High
Threatened Species	Medium	High	High	High	High	High	High	High	Low	Medium	Medium	High
Terrestrial Habitat	Medium	High	High	High	High	High	High	High	Low	Medium	Medium	High
Soils	High	High	High	High	High	High	None	High	Low	Medium	None	None
Agricultural Land	Medium	High	High	High	High	High	None	High	Low	Medium	None	None
Groundwater	None	None	None	None	None	None	None	None	None	None	None	None
Cultural Heritage	None	High	High	High	High	High	High	High	Low	Medium	None	None
Community Capacity	None	None	None	None	None	None	None	None	None	Medium	None	None
ALL REGIONAL ASSETS	Medium	High	High	High	High	High	High	High	Low	Medium	Medium	High

Management Action Priority Matrix

The table below represents the findings of an integration and analysis of the following components:

- the threatening processes priority of action table shown above;
- the management actions from existing management plans shown in Table 21;

- knowledge and evidence supporting past successes (or otherwise) of various management actions in the Mallee or in this Catchment Asset area; and
- the likelihood of implemented management actions addressing or mitigating threatening processes over the life of the RCS.

The criteria used to apply a value to each management action against each Regional Asset is contained in Appendix 6. *Prioritising Potential Management Actions*. The far right column of the table contains a median 'score' for each the management action priority across the Catchment Asset. These 'scores' are the same as those shown in Table 3.

Table 23: Management Action Priority Summary

	Rivers	Wetlands	Threatened Species	Terrestrial Habitat	Soils	Agricultural Land	Groundwater	Cultural Heritage	Community Capacity	Across All Assets
Pest Plant Control	High	High	High	High	High	High	None	High	None	High
Pest Animal Control	High	High	High	High	High	High	None	High	None	High
Habitat Protection	High	High	High	High	Medium	Low	None	High	None	High
Habitat Restoration	Low	Medium	Medium	Medium	Medium	Low	None	Medium	None	Medium
Revegetation	Low	Low	Low	Low	Low	Low	None	Medium	None	Low
Environmental Watering	High	High	High	High	High	Medium	None	Medium	None	High
Soil Erosion Control	Medium	Low	Medium	Medium	High	High	None	High	None	Medium
Threatened Species Interventions	High	High	High	High	None	None	None	None	None	High
Enhancing Land Management Regimes	Medium	High	High	High	High	High	None	High	None	High
Supporting Human Capacity for NRM	Medium	Medium	Medium	Medium	Medium	Medium	None	Medium	Medium	Medium
Supporting Institutional Capacity for NRM	Medium	Medium	Medium	Medium	Medium	Medium	None	Medium	Low	Medium
Supporting Social Capacity for NRM	Low	Low	Medium	Medium	Low	Medium	None	Medium	Medium	Medium
Institutional Planning for NRM	High	High	Medium	Medium	High	Medium	None	High	Medium	Medium
Community Driven Planning for NRM	Low	Medium	Medium	Medium	Medium	Medium	None	Medium	Low	Medium
Landholder Driven Planning for NRM	Low	Medium	Medium	Medium	High	High	None	High	Medium	Medium

Research to improve knowledge	Medium	High	High	High	High	Medium	None	High	Medium	High
Asset condition monitoring and assessment	High	High	High	High	Medium	Medium	None	High	Low	High

Therefore, the highest priority management actions within this Catchment Asset over the life of the Regional Catchment Strategy should be:

- Pest Plant Control;
- Pest Animal Control;
- Habitat Protection;
- Environmental Watering;
- Threatened Species Interventions;
- Enhancing Land Management Regimes;
- Research to improve knowledge; and
- Asset condition monitoring and assessment.

Implementation of these actions should provide the greatest impact on the mitigation of the most concerning threatening processes within this Catchment Asset and make a significant contribution to the achievement of the expected outcomes of the Mallee Regional Catchment Strategy.

Mallee Catchment
Management Authority
www.malleecma.vic.gov.au
PO Box 5017 Mildura 3502
Telephone 03 5051 4377
Facsimile 03 5051 4379